

Datasheet for 200-4233

Ovalbumin Antibody Fluorescein Conjugated**Overview**

Description:	Anti-Ovalbumin (Hen Egg White) (RABBIT) Antibody Fluorescein Conjugated (BULK ORDER) - 200-4233
Item No.:	200-4233
Size:	10 mg
Applications:	Dot Blot, ELISA, WB, FC, IF
Reactivity:	Chicken
Host Species:	Rabbit

Product Details

Background:	Anti-Ovalbumin Antibody detects Ovalbumin. Ovalbumin is the main protein found in egg white, making up 60-65% of the total protein. It is a non-inhibitory serpin. It is the storage protein of egg white. Anti-Ovalbumin Antibody is ideal for investigators involved in Cell Signaling, Immunology and Signal Transduction research.
Synonyms:	rabbit anti-Ovalbumin Antibody FITC Conjugation, fluorescein conjugated rabbit anti-Ovalbumin antibody, Egg albumin antibody, Plakalbumin, Allergen Gal d II, Allergen=Gal d 2, Hen Egg White
Host Species:	Rabbit
Conjugate:	Fluorescein (FITC)
Clonality:	Polyclonal
Format:	IgG
F/P Ratio:	3.6

Target Details

Gene Name:	SERPINB14
Reactivity:	Chicken
Immunogen Type:	Native Protein
Immunogen:	Ovalbumin [Hen Egg White]

Purity/Specificity: Ovalbumin Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Rabbit Serum as well as purified and partially purified Ovalbumin [Hen Egg White]. Cross reactivity against Ovalbumin from other tissues and species may occur but have not been specifically determined.

Relevant Links:

- [200-4233 SDS](#)
- [UniProtKB - P01012](#)
- [NCBI - P01012.2](#)
- [GenelD - 396058](#)

Application Details

Tested Applications: Dot Blot, ELISA, WB

Suggested Applications: FC, IF (Based on references)

Application Note: Anti-Ovalbumin fluorescein conjugated antibody has been tested by ELISA, dot blot, and western blot and is suitable for fluorescent western blot, immunofluorescence microscopy, and flow cytometry or FACS analysis as well as other antibody based fluorescent assays. Researchers should determine optimal titers for applications that are not stated below.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:5,000 - 1:20,000

IF: 1:500 - 1:2,500

IP: 1:100

WB: 1:2,000 - 1:10,000

Formulation

Physical State: Lyophilized

Concentration: 10.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: 0.01% (w/v) Sodium Azide

Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Reconstitution Volume: 1.0 mL

Reconstitution Buffer: Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition: Ambient

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images




Immunofluorescence Microscopy

B, immunohistologic staining. OVA-specific Vα2+ CTLs (red) infiltrated in OVA-expressing tumor tissues (green). Fig 4. PMID: 21628492



Flow Cytometry

j Normalized histogram of singlet-cMyc+ yeast anti-IgE controls stained with secondary reagents alone to assess background signal on yeast, binding to blgE-Fc2-4 alone (100 nM), or co-binding to blgE-Fc2-4 (100 nM) in complex with FcεRIα-Ova (1 μM). Surface bound blgE-Fc2-4 was detected with SA-647 (top) and surface bound FcεRIα-Ova was detected with anti-Ova FITC (bottom). Fig 2. PMID: 34862384

**Western Blot**

Western Blot of Fluorescein conjugated Rabbit anti-Ovalbumin (Hen Egg white) antibody. Lane 1: Non-Reduced Ovalbumin. Lane 2: Opal Pre-stained Ladder (MB-210-0500). Lane 3: Reduced Ovalbumin. Load: 0.05 μ g per lane. Primary antibody: none. Secondary antibody: Fluorescein Rabbit anti-Ovalbumin antibody at 1:1,000 for 60 min at RT. Predicted/Observed size: ~35 kDa.

**Dot Blot**

Dot Blot of Rabbit anti-Ovalbumin (Hen Egg White) Fluorescein Conjugated. Antigen: Ovalbumin. Load: 100 ng, 33.3 ng, 11.1 ng, 3.7 ng, 1.23 ng as indicated. Primary antibody: N/A Secondary antibody: Rabbit anti-Ovalbumin (Hen Egg White) antibody Fluorescein Conjugated 1:1,000 for 60 min at RT. Block: Blocking Buffer for Fluorescent Western Blotting (MB-070) for 1 hour at RT.

References

- Arenas-Hernandez M et al. Specific innate immune cells uptake fetal antigen and display homeostatic phenotypes in the maternal circulation. *J Leukoc Biol.* (2022)
- Pennington LF et al. Directed evolution of and structural insights into antibody-mediated disruption of a stable receptor-ligand complex. *Nat Commun.* (2021)
- Lei F et al. In vivo programming of tumor antigen-specific T lymphocytes from pluripotent stem cells to promote cancer immunosurveillance. *Cancer Res.* (2011)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.