

Datasheet for 200-4372**Lysozyme Antibody Peroxidase Conjugated****Overview**

Description:	Anti-Lysozyme (Hen Egg White) (RABBIT) Antibody Peroxidase Conjugated (BULK ORDER) - 200-4372
Item No.:	200-4372
Size:	20 mg
Applications:	ELISA, WB
Reactivity:	Chicken
Host Species:	Rabbit

Product Details

Background:	Anti-Lysozyme Antibody detects lysozyme. Lysozyme belongs to the glycoside hydrolases enzymes that damage bacterial cell walls by catalyzing hydrolysis of 1,4-beta-linkages between N-acetylmuramic acid and N-acetyl-D-glucosamine residues in a peptidoglycan and between N-acetyl-D-glucosamine residues in chitodextrins. Lysozyme is abundant in a number of secretions, such as tears, saliva, human milk, and mucus. It is also present in cytoplasmic granules of the polymorphonuclear neutrophils. Large amounts of lysozyme can be found in egg white. Anti-Lysozyme Antibody is ideal for investigators involved in Cell Signaling, Neuroscience and Signal Transduction research.
Synonyms:	rabbit anti-Lysozyme Antibody Peroxidase Conjugation, HRP Conjugated rabbit anti-Lysozyme Antibody, 1 4 beta n acetylmuramidase c antibody, EC 3.2.1.17 antibody, LYZ antibody, Lysozyme C antibody, Lysozyme C precursor antibody, LZM antibody, Hen Egg White
Host Species:	Rabbit
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	LYZ
Reactivity:	Chicken

Immunogen Type:	Native Protein
Immunogen:	Anti-Lysozyme Antibody was produced by repeated immunizations with hen egg white Lysozyme protein.
Purity/Specificity:	Anti-Lysozyme Antibody is an IgG fraction purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum as well as purified and partially purified Lysozyme [Hen Egg White]. Cross reactivity against Lysozyme from other tissues and species may occur but have not been specifically determined.
Relevant Links:	• UniProtKB - P00698 • NCBI - NP_990612.1 • GeneID - 396218

Application Details

Tested Applications:	ELISA, WB
Application Note:	Anti-Lysozyme Peroxidase Conjugate Antibody has been tested by western blotting and ELISA and is suitable for IP. Researchers should determine optimal titers for applications that are not stated below.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000
IHC:	1:500 - 1:2,500
IP:	1:100
WB:	1:1,000 - 1:5,000

Formulation

Physical State:	Lyophilized
Concentration:	10 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	2.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition: Ambient

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



ELISA

Release of lysozyme from lysozyme - PCPP complexes (5.7:1 mol/mol) in the (a) absence and (b) presence of HSA as studied in Franz glass diffusion cells - schematically illustrated below (1 mg/mL lysozyme; 10 mg/mL PCPP; 35 mg/mL HSA; PBS, pH 7.4; 2 mL donor compartment; 10 mL acceptor compartment; 50 kDa molecular weight cut off cellulose membrane; release profiles of lysozyme from the donor compartment in the absence of PCPP are shown as diamonds and triangles; two independent experiments were carried out for each dataset with ELISA readouts taken in triplicate, error bars show standard deviation). Fig 7. PMID: 38206583

References

- Hlushko R et al. Macromolecular dimensions of a synthetic polyelectrolyte as a factor in its interactions with protein and cells: desirability for longer chains. *J Mater Chem B*. (2026)
- Marin A et al. Immunopotentiating Polyphosphazene Delivery Systems: Supramolecular Self-Assembly and Stability in the Presence of Plasma Proteins. *Mol Pharm*. (2024)
- Lueckheide M et al. Monitoring Protein Complexation with Polyphosphazene Polyelectrolyte Using Automated Dynamic Light Scattering Titration and Asymmetric Flow Field Flow Fractionation and Protein Recognition Immunoassay. *ACS Polym Au*. (2023)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.