

Datasheet for 200-901-FM9**Transthyretin Antibody****Overview**

Description:	Anti-Transthyretin (CHICKEN) Antibody - 200-901-FM9
Item No.:	200-901-FM9
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse, Rat
Host Species:	Chicken

Product Details

Background:	Transthyretin is a tetrameric carrier protein that transports thyroid hormones in the plasma and cerebrospinal fluid, and retinol (vitamin A) in the plasma. More than 80 different mutations in this gene have been reported; most mutations are related to amyloid deposition, affecting predominantly peripheral nerve and/or the heart. The diseases caused by mutations include familial amyloidotic polyneuropathy, euthyroid hyperthyroxinemia, amyloidotic vitreous opacities, cardiomyopathy, oculoleptomeningeal amyloidosis, meningocerebrovascular amyloidosis, and carpal tunnel syndrome. It has also been suggested that Transthyretin plays an important role in the maintenance of normal cognitive processes during aging, neuropeptide processing and nerve regeneration. It has also been linked to several pathological conditions including Parkinson's disease, schizophrenia, and depression.
Synonyms:	Transthyretin Antibody, CTS, CTS1, PALB, TBPA, HEL111, HsT2651, Transthyretin, ATTR
Host Species:	Chicken
Clonality:	Polyclonal
Format:	IgY

Target Details

Gene Name:	TTR
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-Transthyretin antibody was prepared from chicken egg fractions produced by repeated immunizations with a 17 amino acid synthetic peptide near the internal region of human Transthyretin.
Purity/Specificity:	Anti-Transthyretin Antibody is an IgY fraction that has been affinity purified by immunoaffinity purification. Cross reactivity with Transthyretin from other sources has not been determined.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P02766• GeneID - 7276• NCBI - NP_000362.1

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-Transthyretin Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 16 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000-1:20,000
IF:	20 µg/mL
IHC:	2.5 µg/mL
WB:	1-2 µg/mL

Formulation


Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Wet Ice
----------------------------	---------

Storage Condition:	Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.
Expiration:	Expiration date is one (1) year from date of receipt.


Images



Immunohistochemistry

Spontaneous iPSC-EC differentiation, kidney coculture iPSC-EC differentiation, and previously published iPSC-EC differentiation protocol. (A–D) Spontaneously differentiated iPSC-derived ECs express EC markers (A) VE-Cadherin and (B) ZO-1, a small amount of (C) TTR, but no (D) CA4. (E–H) iPSC-derived ECs differentiated in coculture with primary mouse kidney ECs also express (E) VE-Cadherin, (F) ZO-1, and (G) TTR, but not (H) CA4. (I–L) Likewise, iPSC-derived ECs differentiated using a previously published iPSC-EC differentiation protocol from Rufaihah et al.¹⁸ also express (I) VE-Cadherin, (J) ZO-1, and (K) TTR, but not (L) CA4. DAPI = blue. Scale bars: 100 μm.


Fig 2. PMID: 26720480



Immunohistochemistry


Differentiating choroidal-like ECs via co-culture with RF6A CECs. (A) Schematic illustrating differentiation paradigm: red cells = RF6A CECs, green cells = differentiated iPSC-ECs. (B) GFP expression in live iPSC-L2-RF6A ECs. (C) Carbonic anhydrase IV (CA4; red), GFP (green), and DAPI (blue) expression in iPSC-L2-RF/6A ECs; inset of only CA4 and DAPI expression. (D) GFP (green), TTR (red), and DAPI (blue) expression in iPSC-L2-RF/6A ECs, inset of only TTR and DAPI expression. (E) TTR (green), VE-Cadherin (red), and DAPI (blue) expression in iPSC-L2-RF/6A ECs; inset of only TTR and VE-Cadherin expression; pseudocoloring performed using Fiji. (F) Green fluorescent protein (green) and ZO-1 (red) expression in iPSC-L2-RF/6A ECs; inset of only ZO-1 expression. (G) FOXA2 (gray), VE-Cadherin (red), and DAPI (blue) expression in iPSC-L2-RF/6A ECs; pseudocoloring performed using Fiji. (H) α-SMA (cyan), NG-2 (red), and DAPI expression in iPSC-L2-RF/6A pericytes; pseudocoloring performed using Fiji. Scale bars: 100 μm. Fig 3.

PMID: 26720480




Immunohistochemistry

Characterization and selection of iPSC derived CEC-like cells. (A–C): Immunocytochemical analysis of iPSC derived CEC-like cells using antibodies targeted against TTR (A, C) CA4 (B, C) and CD31 (B) pre-Zeocin selection. (D–F): Immunocytochemical analysis of iPSC derived CEC-like cells using antibodies targeted against TTR (D, E) CA4 (D), CD31 (E) and CD34 (F) post-Zeocin selection. (G): Transmission electron microscopy analysis of a Zeocin-treated iPSC derived CEC-like capillary tube. (F): Expression of EC- and CEC-specific markers by Zeocin-selected cells and controls, as detected via rt-PCR. (H): Rt-PCR analysis demonstrating expression of the endothelial cell markers CA4, PLVAP, ICAM1, and vWF. (I): Schematic diagram illustrating the steps and timeline required from generating patient specific iPSCs to derivation of Zeocin selected CEC-like cells. All scale bars represent 100 μm. Abbreviations: CEC, choroidal endothelial cell; DAPI, 4',6-diamidino-2-phenylindole; EB, embryoid body; iPSC, induced pluripotent stem cell; TTR, transthyretin. Fig 4. PMID: 28474838



Immunohistochemistry


Immunohistochemistry of Transthyretin antibody. Tissue: human lung tissue. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: Transthyretin antibody at 2.5 μg/mL for 1 h at RT. Secondary antibody: Peroxidase chicken secondary antibody at 1:10,000 for 45 min at RT. Localization: Transthyretin is nuclear. Staining: Transthyretin as precipitated light brown signal with brown nuclear counterstain.



Immunofluorescence Microscopy

Immunofluorescence Microscopy of Transthyretin antibody.

Tissue: human lung tissue. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: Transthyretin antibody at 20 $\mu\text{g}/\text{mL}$ for 1 h at RT. Secondary antibody: Fluorescein chicken secondary antibody at 1:10,000 for 45 min at RT. Localization: Transthyretin is nuclear. Staining: Transthyretin as red fluorescent signal.



Western Blot

Western Blot of Transthyretin antibody. Lane 1: Human lung tissue lysate with Transthyretin antibody at 1 $\mu\text{g}/\text{mL}$. Lane 2: Human lung tissue lysate with Transthyretin antibody at 2 $\mu\text{g}/\text{mL}$. Load: 35 μg per lane. Secondary antibody: Peroxidase chicken secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 16 kDa, 25 kDa for Transthyretin. Other band(s): Transthyretin splice variants and isoforms.

References

- Songstad et al. Connective Tissue Growth Factor Promotes Efficient Generation of Human Induced Pluripotent Stem Cell-Derived Choroidal Endothelium. *STEM CELLS Translational Medicine* (2017)
- Songstad et al. Generating iPSC-Derived Choroidal Endothelial Cells to Study Age-Related Macular Degeneration. *Investigative Ophthalmology & Visual Science* (2015)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.