

**Datasheet for 200-B26-N88****CD69 Allophycocyanin Antibody****Overview**

<b>Description:</b>	Anti-CD69 (Armenian Hamster) Allophycocyanin Conjugated Monoclonal Antibody - 200-B26-N88
<b>Item No.:</b>	200-B26-N88
<b>Size:</b>	100 µg
<b>Applications:</b>	FC
<b>Reactivity:</b>	Mouse
<b>Host Species:</b>	Hamster Armenian

**Product Details**

<b>Background:</b>	CD69 is a 60 kD type II membrane protein composed of a 27/33 kD disulfide-linked homodimer, also known as Very Early Activation Antigen (VEA), AIM, EA1, MLR3, and gp34/28. It is expressed on a subset of thymocytes and platelets. CD69 is rapidly induced on activated T and B cells, neutrophils, and NK cells. It is a C-type lectin, closely related to the NKR-P1 and Ly-49 NK cell activation molecules. CD69 is involved in the early events of cell activation and thymocyte positive selection.
<b>Synonyms:</b>	Early activation antigen CD69, CD69, Cd69
<b>Host Species:</b>	Hamster Armenian
<b>Conjugate:</b>	Allophycocyanine (APC)
<b>Clonality:</b>	Monoclonal
<b>Clone ID:</b>	H1.2F3
<b>Format:</b>	IgG
<b>F/P Ratio:</b>	1-2

**Target Details**

<b>Gene Name:</b>	Cd69
<b>Reactivity:</b>	Mouse

<b>Immunogen Type:</b>	Other
<b>Immunogen:</b>	Anti-CD69 Antibody (Monoclonal) was produced by repeated immunizations with Mouse dendritic epidermal T cell line Y245.
<b>Purity/Specificity:</b>	Allophycocyanin conjugated CD69 Monoclonal Antibody was purified from tissue culture supernatant via affinity chromatography and is directed against mouse CD69. Cross reactivity with CD69 from other sources has not been tested. Anti-CD69 is conjugated with APC under optimal conditions and the solution is free of unconjugated APC.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P37217</a></li><li>• <a href="#">NCBI - NP_001028294.1</a></li><li>• <a href="#">GeneID - 12515</a></li></ul>

## Application Details

<b>Tested Applications:</b>	FC
<b>Application Note:</b>	Anti-CD69 is tested for FLOW and useful for Immunohistochemistry and Immunoprecipitation using mouse spleen cells, or an appropriate cell type (where indicated). Researchers should determine optimal titers for applications that are not stated.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>FC:</b>	10 µL/10 <sup>6</sup> cells (0.1 µg)
<b>IHC:</b>	User Optimized
<b>IP:</b>	User Optimized

## Formulation


<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	0.2mg/mL Sufficient to run approximately 100 tests
<b>Buffer:</b>	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.09% (w/v) Sodium Azide
<b>Stabilizer:</b>	None

## Shipping & Handling

<b>Shipping Condition:</b>	Wet Ice
----------------------------	---------

<b>Storage Condition:</b>	Store vial at 4° C prior to opening. Dilute only prior to immediate use. This product is stable at 4° C as an undiluted liquid. Use subdued lighting during handling and incubation of cells prior to analysis. Store reagent in the dark. DO NOT FREEZE.
<b>Expiration:</b>	Expiration date is six (6) months from date of receipt.

## Images



### Flow Cytometry

Flow Cytometry of anti-CD69 Allophycocyanin Conjugated Monoclonal Antibody. Cells: C57Bl/6 mouse splenocytes. Stimulation: Con-A (2 days). Antibody: (Unfilled Line) APC armenian hamster IgG isotype control; (Filled Line) Allophycocyanin Anti-CD69 antibody.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.