

Datasheet for 209-301-B31S**IL-17F Antibody****Overview**

Description:	Anti-Human IL-17F (MOUSE) Monoclonal Antibody - 209-301-B31S
Item No.:	209-301-B31S
Size:	25 µL
Applications:	IHC, WB
Reactivity:	Human
Host Species:	Mouse

Product Details

Background: Anti-L-17F recognizes IL-17F (also known as Cytokine ML-1 or Interleukin-24). IL-17F is produced and secreted by CD8+ T cells, NK cells, NKT cells and LT α i cells. The main functions of IL-17F are neutrophil recruitment and immunity to extracellular pathogen. More importantly, IL-17F drives inflammation and auto-immunity. IL-17A and IL-17F are by far the best characterized cytokines of the IL-17 cytokine family. IL-17F dimerizes in a parallel fashion similar to nerve growth factor and other neurotrophins. Its dimerization is critical to fulfill its activity. When secreted by activated T cells, IL-17F can stimulate the production of other cytokines such as IL-6, IL-8 granulocyte colony-stimulating factor and, can stimulate cartilage matrix turnover. Defects in IL17F are the cause of familial candidiasis type 6 (CANDF6). CANDF6 is a rare disorder with altered immune responses and impaired clearance of fungal infections, selective against Candida. Anti-IL-17E cytokine antibody is ideal for investigators involved in Immunology, Signal Transduction research, Cancer and Inflammatory pathologies.

Synonyms:	mouse anti-IL-17F antibody, mouse anti-Interleukin-17F antibody, Cytokine ML-1, Interleukin-24
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	4H1629.1
Format:	IgG1

Target Details

Gene Name:	IL17F
-------------------	-------

Reactivity:	Human
Immunogen Type:	Recombinant Protein
Immunogen:	Anti-IL-17F (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with mature full length recombinant human IL-17F produced in E.coli followed by hybridoma development.
Purity/Specificity:	Anti-Human IL-17F (MOUSE) Monoclonal Antibody was purified from concentrated tissue culture supernate by Protein G chromatography followed by extensive dialysis against the buffer stated above. This antibody is specific for human IL-17F protein. A BLAST analysis was used to suggest cross-reactivity with IL-17F from human sources based on 100% homology with the immunizing sequence. Cross-reactivity with IL-17F from other sources has not been determined.
Relevant Links:	• UniProtKB - Q96PD4 • NCBI - NP_443104.1 • GeneID - 112744

Application Details

Tested Applications:	IHC, WB
Application Note:	Anti-Human IL-17F antibody has been tested for use in IHC and Western Blot. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:50,000
IHC:	5µg/mL
WB:	0.5µg/mL


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling


Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Immunohistochemistry

Immunohistochemistry of Mouse Anti-IL-17F antibody. Tissue: human colon tissue. Fixation: formalin-fixed, paraffin-embedded. Primary antibody: isotype control (top left), Mouse Anti-IL-17F antibody (full right and bottom left) at 5 ug/ml.



Western Blot

Western Blot of Mouse Anti-IL-17F antibody. Lane 1: human full length recombinant IL-17F protein. Lane 2: mouse full length recombinant IL-17F protein. Lane 3: rat full length recombinant IL-17F protein. Load: 20 ng/lane. Primary antibody: Anti-IL-17F antibody at 0.5ug/mL for overnight at 4°C.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.