

Datasheet for 209-401-B50S**IL-33 Antibody****Overview**

Description:	Anti-Human IL-33 (RABBIT) Antibody - 209-401-B50S
Item No.:	209-401-B50S
Size:	25 µL
Applications:	WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	IL-33 (also known as Interleukin-33, Interleukin-1 family member 11, IL-1F11, nuclear factor from high endothelial venules and NF-HEV) is a cytokine that binds to and signals through IL1RL1/ST2 and its stimulation recruits MYD88, IRAK1, IRAK4, and TRAF6, followed by phosphorylation of MAPK3/ERK1 and/or MAPK1/ERK2, MAPK14, and MAPK8. IL-33 induces T helper type 2-associated cytokines. IL-33 is a secreted cytokine that is expressed at high levels in high endothelial venules found in tonsils, Peyer patches and mesenteric lymph nodes and is almost undetectable in placenta. The 31 kDa precursor is proteolytically converted to an 18 kDa mature form by CASP1. Anti-IL-33 antibody is ideal for investigators involved in Cardiovascular and Immunology research.
Synonyms:	rabbit anti-Interleukin-33 antibody, rabbit anti-IL-33 antibody, Interleukin-1 family member 11, IL-1F11, nuclear factor from high endothelial venules and NF-HEV
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	IL33
Reactivity:	Human
Immunogen Type:	Recombinant Protein

Immunogen:	This purified antibody was prepared from whole rabbit serum produced by repeated immunizations with full length recombinant human IL-33 protein.
Purity/Specificity:	This purified antibody has been heated to 56°C for 30 minutes. In ELISA and other immunoreactive assays, this antibody will recognize both native and recombinant human IL-33 in cell supernatants and certain body fluids. A control of similarly diluted normal rabbit IgG is recommended.
Relevant Links:	• NCBI - 15559209 • UniProtKB - O95760 • GeneID - 90865

Application Details

Tested Applications:	WB
Application Note:	This purified antibody has been tested in western blotting. Reactivity is also expected in ELISA, neutralizations, radioimmunoassay and immunohistochemistry. The endotoxin content is estimated to be <10 pg/μl by the LAL method. By western blot a band approximately 18 kDa in size corresponding to mature human IL-32α protein is expected in the appropriate cell lysate or extract. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000-1:100,000
Neutralization:	1:400
WB:	1:500-1:2,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/ml by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	None
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

Expiration: Expiration date is three (3) months from date of receipt.

Images



Western Blot

Western blot using Rockland's anti-Human IL-33 antibody shows detection of a band ~18 kDa in size corresponding to recombinant human IL-33 (lane 1). The identity of the higher molecular weight band is unknown. Molecular weight markers are also shown (M). After transfer, the membrane was blocked overnight with 3% BSA in TBS followed by reaction with primary antibody at a 1:1,000 dilution. Detection occurred using peroxidase conjugated anti-Rabbit IgG (p/n 611-103-122) secondary antibody diluted 1:40,000 in blocking buffer (p/n MB-070) for 30 min at RT followed by reaction with FemtoMax™ chemiluminescent substrate. Image was captured using VersaDoc™ MP 4000 imaging system (Bio-Rad).

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.