

Datasheet for 209-403-310**IL-6 Antibody Peroxidase Conjugated****Overview**

Description:	Anti-Human IL-6 (RABBIT) Antibody Peroxidase Conjugated - 209-403-310
Item No.:	209-403-310
Size:	100 µg
Applications:	WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	IL-6 is a secreted cytokine with a wide variety of biological functions. It is a potent inducer of the acute phase response and plays an essential role in the final differentiation of B-cells into Ig-secreting cells. Involved in lymphocyte and monocyte differentiation. IL-6 induces myeloma and plasmacytoma growth and induces nerve cells differentiation and acts on B-cells, T-cells, hepatocytes, hematopoietic progenitor cells and cells of the CNS. IL-6 also acts as a myokine. It is discharged into the bloodstream after muscle contraction and acts to increase the breakdown of fats and to improve insulin resistance.
Synonyms:	rabbit Anti-IL-6 peroxidase conjugated antibody, rabbit anti-interleukin-6 peroxidase conjugated antibody, HSF, Hybridoma growth factor, Hybridoma plasmacytoma growth factor, IFNB2, IL 6, IL6 protein, Interferon beta 2, Interleukin 6, BSF2, CDF
Host Species:	Rabbit
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	IL6
Reactivity:	Human
Immunogen Type:	Recombinant Protein

Immunogen:	This purified antibody was prepared from whole rabbit serum produced by repeated immunizations with full length recombinant human IL-6 produced in E.coli.
Purity/Specificity:	This purified antibody has been heated to 56°C for 30 minutes. In ELISA and other immunoreactive assays, this antibody will recognize both native and recombinant human IL-6 in cell supernatants and certain body fluids. In Western blot analysis of natural cell products or human body fluids, multiple bands of IL-6 will appear due to the variable amount of glycosylation on the molecule. A control of similarly diluted normal rabbit IgG is recommended. This product has minimal reactivity with mouse IL-6.
Relevant Links:	• UniProtKB - P05231 • NCBI - NP_000591.1 • GenelD - 3569

Application Details

Tested Applications:	WB
Application Note:	This purified antibody has been tested in western blotting. Reactivity is also expected in ELISA, neutralizations, radioimmunoassay and immunohistochemistry. The endotoxin content is estimated to be <10 pg/μl by the LAL method. By western blot a band approximately 23.7 kDa in size corresponding to native human IL-6 protein is expected in the appropriate cell lysate or extract. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000-1:50,000
IHC:	1:500-1:2,500
WB:	1:1,000-1:5,000


Formulation

Physical State:	Lyophilized
Concentration:	1.1 mg/ml by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	100 μL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling


Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Rabbit Anti-IL-6 Antibody Peroxidase Conjugated. Lane 1: Opal Prestained Molecular Weight Marker (p/n MB-210-0500). Lane 2: IL-6/MOLT-4 lysate (p/n W09-001-GK2) [0.01/10µg]. Lane 3: IL-6/MOLT-4 lysate (p/n W09-001-GK2) [0.005/10µg]. Lane 4: IL-6/MOLT-4 lysate (p/n W09-001-GK2) [0.002/10µg]. Primary Antibody: Anti-IL-6 HRP conjugated at 1.0µg/mL overnight at 2-8°C. Secondary Antibody: Goat Anti-Rabbit IgG HRP mx10 (p/n 611-103-122) at 1:70,000 for 30min at RT. Blocking: BlockOut Buffer (p/n MB-073) at RT for 1hr. Predicted/Observed MW: ~20.6kDa. Exposure: 0.5sec.



Western Blot

Western Blot of Peroxidase conjugated Rabbit anti-Human IL-6 antibody. Lane 1: Human IL-6. Lane 2: none. Load: 50 ng per lane. Primary antibody: none. Secondary antibody: Peroxidase Human IL-6 secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 20 kDa for Human IL-6. Other band(s): none.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.