

Datasheet for 210-301-C56S**IL-35 EBI3 Antibody****Overview**

Description:	Anti-Mouse EBI-3 (MOUSE) Monoclonal Antibody - 210-301-C56S
Item No.:	210-301-C56S
Size:	25 µL
Applications:	Neutralization
Reactivity:	Mouse
Host Species:	Mouse

Product Details

Background: The cytokine Interleukin 27 (IL-27) is produced in response to inflammation. It is made by activated antigen presenting cells including monocytes, endothelial cells, and dendritic cells. IL-27 consists of a heterodimeric combination of Epstein-Barr virus-induced molecule 3 (EBI3, or IL-27B) non-covalently linked with IL-27 p28 (or IL-27A). It is a regulator of T helper cell development and suppressor of T-cell proliferation. IL-27 has both pro- and anti-inflammatory properties. It can stimulate cytotoxic T cell activity and induce isotype switching in B-cells. It has diverse effects on innate immune cells. It induces monocytes and mast cells to secrete pro-inflammatory cytokines. When infection is present, IL-27 induces naive CD4+ T cells to proliferate and develop Th1 cell responses. As an anti-inflammatory regulator, IL-27 can inhibit Th1 or Th2 responses and restrict the strength and duration of adaptive immune responses.

The IL-27 p28 subunit, a 28 kDa glycoprotein belonging to the type I cytokine family, is homologous to IL-12 p35, IL-23 p19, and IL-6. The EBI3 (Epstein-Barr virus-induced molecule 3, or IL-27B) subunit is a 34 kDa glycoprotein containing two fibronectin type III domains, and belongs to the type I cytokine receptor family. It can exist as a homodimer and can also heterodimerize with IL-27 p28 or IL-12 p35 subunit. It is homologous to the p40 subunit of IL-12 and IL-23 and to the extracellular domain of IL-6 R.

Synonyms:	mouse anti-EBI3 antibody, mouse anti-IL-35 antibody, Epstein-Barr virus induced 3 protein, Interleukin-27 subunit beta, IL-27 subunit beta, IL-27B, Epstein-Barr virus-induced gene 3 protein, EBV-induced gene 3 protein, EBI-3, EBI3, IL-35, IL35
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	V1.4F5.25

Format: IgG2b

Target Details

Gene Name: Ebi3

Reactivity: Mouse

Immunogen Type: Recombinant Protein

Immunogen: Anti-EBI-3 (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with mature full length recombinant mouse EBI-3 produced in E.coli followed by hybridoma development.

Purity/Specificity: Anti-EBI-3 was purified from concentrated tissue culture supernate by Protein A chromatography followed by extensive dialysis against the buffer stated above. This antibody is specific for mouse EBI3 protein. A BLAST analysis was used to suggest cross-reactivity with EBI3 from mouse sources based on 100% homology with the immunizing sequence. Cross-reactivity with EBI-3 from other sources has not been determined.

Relevant Links:

- [GenelD - 50498](#)
- [NCBI - NP_056581.1](#)
- [UniProtKB - O35228](#)

Application Details

Tested Applications: Neutralization

Application Note: EBI-3 antibody has been tested for use in IP and Neutralization. Specific conditions for reactivity should be optimized by the end user.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:10,000

IP: 1µg/mL

Neutralization: 10µg/mL

WB: 1:1000

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: 0.01% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Neutralization

Neutralization: Murine CD4+CD25- (T_{conv}) were purified by FACS, activated with anti-CD3/CD28-coated beads in the presence of a titration of mIL-35 and 10µg/ml of 210-301-C56 antibody. Percent suppression was determined by calculating the percent decrease in counts per minute of activated T_{conv} plus or minus murine IL-35.



Neutralization

Neutralization: Human CD4+CD25- (Tconv) were purified by FACS from cord blood, activated with anti-CD3/CD28-coated beads and 10IU IL-2 in the presence of hIL-35 and a titration of an isotype control or 210-301-C56 antibody. Percent neutralization was determined by calculating the percent decrease in counts per minute of activated Tconv plus (0%) or minus (100%) human IL-35 in the presence and absence of Ab.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.