

Datasheet for 210-401-B66**IL-35 EBI3 Antibody****Overview**

Description:	Anti-Mouse EBI-3 (RABBIT) Antibody - 210-401-B66
Item No.:	210-401-B66
Size:	100 µg
Applications:	WB
Reactivity:	Mouse, Rat
Host Species:	Rabbit

Product Details

Background: The cytokine Interleukin 27 (IL-27) is produced in response to inflammation. It is made by activated antigen presenting cells including monocytes, endothelial cells, and dendritic cells. IL-27 consists of a heterodimeric combination of Epstein-Barr virus-induced molecule 3 (EBI3, or IL-27B) non-covalently linked with IL-27 p28 (or IL-27A). It is a regulator of T helper cell development and suppressor of T-cell proliferation. IL-27 has both pro- and anti-inflammatory properties. It can stimulate cytotoxic T cell activity and induce isotype switching in B-cells. It has diverse effects on innate immune cells. It induces monocytes and mast cells to secrete pro-inflammatory cytokines. When infection is present, IL-27 induces naive CD4+ T cells to proliferate and develop Th1 cell responses. As an anti-inflammatory regulator, IL-27 can inhibit Th1 or Th2 responses and restrict the strength and duration of adaptive immune responses.

The IL-27 p28 subunit, a 28 kDa glycoprotein belonging to the type I cytokine family, is homologous to IL-12 p35, IL-23 p19, and IL-6. The EBI3 (Epstein-Barr virus-induced molecule 3, or IL-27B) subunit is a 34 kDa glycoprotein containing two fibronectin type III domains, and belongs to the type I cytokine receptor family. It can exist as a homodimer and can also heterodimerize with IL-12 p35. It is homologous to the p40 subunit of IL-12 and IL-23 and to the extracellular domain of IL-6 R. EBI3 can heterodimerize also with IL-12 p35, or can exist as a homodimer.

Synonyms: rabbit anti-EBI3 antibody, rabbit anti-IL-35 antibody, Epstein-Barr virus induced 3 protein, Interleukin-27 subunit beta, IL-27 subunit beta, IL-27B, Epstein-Barr virus-induced gene 3 protein, EBV-induced gene 3 protein, EBI-3, EBI3, IL-35, IL35

Host Species: Rabbit

Clonality: Polyclonal

Format: IgG

Target Details

Gene Name: Ebi3

Reactivity: Mouse, Rat

Immunogen Type: Recombinant Protein

Immunogen: This antibody was prepared by repeated immunizations with recombinant mouse EBI3.

Purity/Specificity: This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. This antibody is specific for mouse and rat EBI-3 protein. Cross reactivity does NOT occur with human EBI-3. Cross-reactivity with IL-27 from other sources has not been determined.

Relevant Links:

- [UniProtKB - O35228](#)
- [NCBI - NP_056581](#)
- [GenelD - 50498](#)

Application Details

Tested Applications: WB

Application Note: IL-27 is expressed in activated antigen presenting cells including monocytes, endothelial cells, and dendritic cells, for example mouse CD4 splenocytes. This purified antibody has been tested for use in western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 23 kDa in size corresponding to the mature mouse EBI3 protein, a non-glycosylated polypeptide chain consisting of 207 amino acids, by western blotting in appropriate cell lysate or extract.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:1,000 - 1:5,000

WB: 1:500 - 1:2,000

Formulation

Physical State: Lyophilized

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: None

Stabilizer: None

Reconstitution Volume: 100 μ L

Reconstitution Buffer: Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition: Ambient

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is six (6) months from date of receipt.

Images



Western Blot

Recombinant mouse EBI-3 was loaded on to an SDS-PAGE gel at 0.1 μ g and after separation, transferred to nitrocellulose. The expected band is a non-glycosylated polypeptide chain consisting of 207 amino acids and is approximately 23 kDa in size. The membrane was blocked with 1% BSA in TBST for 30 min at RT, followed by incubation with primary antibody diluted 1:1,000 in 1% BSA in TBST overnight at 4°C. After washes, the blot was reacted with secondary antibody HRP Goat anti-Rabbit IgG antibody (p/n 611-103-122) diluted 1:40,000 in blocking buffer (p/n MB-070) for 30 min at RT. Data was collected using Bio-Rad VersaDoc® 4000 MP imaging system.

**Western Blot**

Western Blot of Rabbit Anti-Mouse EBI3 Antibody. Lane 1: Molecular Weight. Lane 2: Mouse EBI-3. Primary Antibody: Rabbit Anti-Mouse EBI3 at 1:1000 overnight at 2-8°C. Secondary Antibody: Rabbit anti-Mouse IgG peroxidase (p/n 610-4302) at 1:40000 for 30mins at RT. Block: (p/n MB-070).

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.