

Datasheet for 600-101-387**R-Phycoerythrin Antibody****Overview**

Description:	Anti-R-Phycoerythrin (GOAT) Antibody - 600-101-387
Item No.:	600-101-387
Size:	1 mg
Applications:	ELISA, IHC
Reactivity:	Phycoerythrin
Host Species:	Goat

Product Details

Background:	Anti-R-Phycoerythrin Antibody generated in goat recognizes RPE. Phycoerythrin (PE) is a red protein-pigment present in red algae and cryptophytes and an accessory pigment to the main chlorophyll pigments responsible for photosynthesis. R-phycoerythrin is predominantly produced by red algae. The molecular weight of R-PE is 240kDa. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
Synonyms:	goat anti R-Phycoerythrin Antibody, goat anti-R-Phycoerythrin Antibody, goat anti Phycoerythrin
Host Species:	Goat
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Phycoerythrin
Immunogen Type:	Native Protein
Immunogen:	This affinity purified antibody was prepared from whole goat serum produced by repeated immunizations with highly purified R-Phycoerythrin from the seaweed gracilaria.

Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using a R-Phycoerythrin coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum and R-Phycoerythrin conjugated IgG. This antibody will cross react with B-Phycoerythrin. Reactivity with other phycobiliproteins is unknown.
Relevant Links:	• 600-101-387 SDS • UniProtKB - Q7SIG0

Application Details

Tested Applications:	ELISA
Suggested Applications:	IHC (Based on references)
Application Note:	Anti-R-Phycoerythrin Antibody is tested by ELISA and suitable for western blotting, fluorescent assays, immunoprecipitation, immunodiffusion, conjugation to magnetic particles (beads) and most other immunological methods requiring lot-to-lot consistency, high titer and specificity. Although not specifically tested, this reagent should also be useful for FACS and other fluorescent immunoassays. R-Phycoerythrin (240 kDa) is a labile molecule that may dissociate into components upon exposure to reducing or denaturing agents. Reaction with low molecular fragments is typically noted by western blot.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:10,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Wet Ice
----------------------------	---------

Storage Condition:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

References

- Veenstra et al. Diabetic Retinopathy: Retina-Specific Methods for Maintenance of Diabetic Rodents and Evaluation of Vascular Histopathology and Molecular Abnormalities. *Current Protocols in Mouse Biology* (2015)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.