

Datasheet for 600-103-200

GST Antibody HRP Conjugated**Overview**

Description:	Anti-GST (GOAT) Antibody Peroxidase Conjugated - 600-103-200
Item No.:	600-103-200
Size:	1 mg
Applications:	ELISA, WB
Reactivity:	GST-Tag
Host Species:	Goat

Product Details

Background:	GST Antibody HRP Conjugated is specific for the GST affinity tag. Affinity tags are appended to proteins thereby allowing them to be purified from their crude biological source using an affinity technique. Common affinity tags include glutathione-S-transferase (GST), chitin binding protein (CBP), maltose binding protein (MBP), and the poly-Histidine or HIS-tag. Rockland produces a wide range of GST antibodies in our laboratories. Select GST antibodies from several monoclonal and/or polyclonal GST antibodies listed below. Select appropriate GST antibodies for your research by isotype, epitope, applications and species reactivity. GST (Glutathione-S-Transferase) is a protein expression tag commonly used in molecular biology. Anti-GST will react with synthetic construct present in most known GST containing cloning or expression vectors. GST is responsible for the conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. The amino acid sequence GST is highly conserved in most organisms including mammals. GST exists as a 26 kDa homodimer.
Synonyms:	goat anti-GST antibody HRP conjugation, peroxidase conjugated goat anti-GST antibody, Glutathione-S-Transferase, GST antibody, anti-GST antibody, anti-Glutathione-S-Transferase antibody
Host Species:	Goat
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	GST-Tag
Immunogen Type:	Native Protein
Immunogen:	Glutathione-S-Transferase [Schistosoma japonicum]
Purity/Specificity:	Anti-GST HRP Conjugated antibody was prepared from monospecific antiserum by immunoaffinity chromatography using GST coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, anti-Peroxidase, as well as purified and partially purified Glutathione-S-Transferase [Schistosoma japonicum]. Cross reactivity against Glutathione-S-Transferase from other sources may occur but has not been specifically determined.
Relevant Links:	<ul style="list-style-type: none">• 600-103-200-SDS

Application Details

Tested Applications:	ELISA, WB
Application Note:	Anti-GST peroxidase conjugated Antibody has been tested by ELISA and western blot and is suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using peroxidase conjugates requiring lot-to-lot consistency. Optimal concentrations in these or other immunoassays should be determined by the researcher.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000
IHC:	1:500 - 1:2,000
WB:	1:2,000 - 1:5,000

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition: Ambient

Storage Condition: Store anti-GST at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.


Images



ELISA


A, The expression of recombinant NHDL and r18D11 by stably transfected CHO-K1SP cell pools (fed-batch) was compared to transient expression by HEK293E cells (batch-batch). Cell culture supernatants were analyzed for IgG levels using ELISA. Results are given as single value mg IgG per E06 CHO cells (n = 1), and mean mg IgG ± S.E.M. per E06 HEK293E cells for NHDL (n = 3) and r18D11 (n = 6). B, Binding of increasing concentrations of CHO- and HEK293E-produced NHDL or r18D11 to recombinant human C1q or recombinant human Fcγ receptors hFcγRI, hFcγRIIa, and hFcγRIIIa at pH 7.4 was measured by ELISA. As a positive control, a recombinant human IgG1 antibody (bevacizumab (Avastin®), Roche) was included in the analysis. The results are shown as relative units for the absorbance at 450 nm (n = 2; mean ± S.D.). C, Human whole blood was incubated with CHO- or HEK293E-produced r18D11 (1.5, 15 and 30 μg/mL) or NHDL (15 μg/mL) in the presence of 100 ng/mL upLPS. As controls, 15 μg/mL r18D11 or NHDL were incubated in the presence of PBS, only. Plasma IL-6 was measured using Bioplex technology. Data are given as pg/mL (n = 2; mean ± S.D.). Fig 5.

PMID: 31671278



ELISA

Human albumin binds IgA1-fused ABD in a pH-independent manner but human FcRn in a pH-dependent manner. A-B) ELISA showing binding of titrated amounts of ABD-fused IgA1 variants to human albumin at both pH 5.5 and 7.4. The numbers represent the mean±s.d. of duplicates. C-D) ELISA showing pH-dependent binding of ABD-fused IgA1 in complex with human albumin to human FcRn. The numbers represent the mean±s.d. of duplicates. Figure A and C were made with bioRender. Fig 2. PMID: 33691596



Western Blot

Western Blot Results of Goat Anti-GST Antibody Peroxidase Conjugated. Lane 1: rec GST (p/n 000-001-200)/HeLa WCL (p/n W09-000-364) [0.03/10µg]. Lane 2: HeLa Whole Cell Lysate (p/n W09-000-364) [10µg]. Lane 3: 12 Epitope GST Lysate (p/n MB-302-0100) [0.5µg]. Lane 4: Opal Prestained 11-245 kDa Molecular Weight Marker (p/n MB-210-0500). Antibody: Goat Anti-GST Peroxidase Conjugated Antibody at 1.0µg overnight at 4°C. BlockOut blocking buffer (p/n MB-073) for 1hr at RT. Predicted MW: 25kDa. Observed MW: 25kDa in lane 1. 48kDa in lane 4 for 12 Epitope GST Lysate. Exposure: 0.03 seconds.

References

- Bertoglio F et al. Antibodies to coagulase of *Staphylococcus aureus* crossreact to Efb and reveal different binding of shared fibrinogen binding repeats. *Front Immunol.* (2023)
- Grevys A et al. Antibody variable sequences have a pronounced effect on cellular transport and plasma half-life. *iScience* (2022)
- Mester S et al. Extended plasma half-life of albumin-binding domain fused human IgA upon pH-dependent albumin engagement of human FcRn in vitro and in vivo. *Mabs.* (2021)
- Lau C. et al. NHDL, a Recombinant V L/V H Hybrid Antibody Control for IgG2/4 Antibodies. *MABs* (2020)
- Lee, CH et al. An engineered human Fc domain that behaves like a pH-toggle switch for ultra-long circulation persistence. *Nature Communications* (2019)
- Lee et al. IgG Fc domains that bind C1q but not effector Fcγ receptors delineate the importance of complement-mediated effector functions. *Nature Immunology* (2017)
- Higai K et al. NKG2D and CD94 bind to heparin and sulfate-containing polysaccharides. *Biochem Biophys Res Commun.* (2009)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.