

Datasheet for 600-141-215**GFP Antibody DyLight™ 488 Conjugated****Overview**

Description:	Anti-GFP (GOAT) Antibody DyLight™ 488 Conjugated (Min X Hu Ms & Rt Serum Proteins) - 600-141-215
Item No.:	600-141-215
Size:	100 µg
Applications:	Dot Blot, WB, FC, IF, IHC, IP, Multiplex
Reactivity:	GFP, eGFP, rGFP
Host Species:	Goat

Product Details

Background:	GFP DyLight™ 488 Conjugated Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
Synonyms:	goat anti-GFP Antibody DyLight™ 488 Conjugation, DyLight™ 488 conjugated goat anti-GFP antibody, Green Fluorescent Protein, GFP antibody, Green Fluorescent Protein antibody, EGFP, enhanced Green Fluorescent Protein, Aequorea victoria, Jellyfish
Host Species:	Goat
Conjugate:	DyLight™ 488
Clonality:	Polyclonal
Format:	IgG
F/P Ratio:	5.7

Target Details

Reactivity:	GFP, eGFP, rGFP
Immunogen Type:	Recombinant Protein
Immunogen:	The immunogen is a Green Fluorescent Protein (GFP) fusion protein corresponding to the full length amino acid sequence (246aa) derived from the jellyfish Aequorea victoria.

Purity/Specificity: GFP Dylight™ 488 Conjugated Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Green Fluorescent Protein (*Aequorea victoria*) coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum and purified and partially purified Green Fluorescent Protein (*Aequorea victoria*). No reaction was observed against Human, Mouse or Rat serum proteins.

Relevant Links:

- [UniProtKB - P42212](#)

Application Details

Tested Applications:	Dot Blot, WB
Suggested Applications:	FC, IF, IHC, IP, Multiplex (Based on references)
Application Note:	Anti-GFP Dylight™ 488 Conjugated Antibody has been tested by dot blot and western blot. The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000 - 1:40,000
FLISA:	>1:10,000
IF:	>1:5,000
WB:	1:10,000

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	100 µL
Reconstitution Buffer:	Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition: Ambient

Storage Condition: Store Conjugated anti-GFP at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Bot of Goat Anti-GFP (GOAT) Antibody DyLight™ 488 Conjugated.

Lane 1: Opal Prestained Molecular Weight Marker (p/n MB-210-0500).

Lane 2: GFP / HEK293T Whole Cell Lysate Reduced [0.05/10µg].

Lane 3: HEK293T Whole Cell Lysate Reduced (p/n W09-001-GX5) [10µg].

Lane 4: GFP / NIH-3T3 Whole Cell Lysate Reduced [0.05/10µg].

Lane 5: NIH-3T3 Whole Cell Lysate Reduced (p/n W10-000-358) [10µg].


Lane 6: GFP / PC-12 Whole Cell Lysate Reduced [0.05/10µg].

Lane 7: PC-12 Whole Cell Lysate Reduced (p/n W12-001-GL9) [10µg].

Secondary Antibody: Goat Anti-GFP (GOAT) Antibody DyLight™ 488 Conjugated at 1.0µg/mL overnight at 2-8°C.

Blocking Buffer: Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) for 1 hour at RT.

Expect: 27kDa.



Immunofluorescence Microscopy

(A) Immunofluorescence (IF) stainings of CX3CR1hi Macs in ear skin of 10-day- and 8-week-old Cx3cr1+/gfp mice. Scale bars: 100 μ m.

(B) Localization of CX3CR1hi Macs in relation to collagen IV + basement membranes and β 3-tubulin+ nerve bundles in ear skin of a Cx3cr1+/gfp mouse (P24). Insets: direct alignment of CX3CR1hi Macs along nerves. Scale bar: 100 μ m. Yellow fluorescent protein (YFP) and in rare cases (Figure 4B) the endogenous GFP signal in Cx3cr1+/gfp mice, was enhanced with anti-GFP DyLight488 antibody (Rockland).

(C) F4/80-positive staining of nerve-associated CX3CR1hi cells (arrowheads). Scale bars: 20 μ m.

(D) Confocal microscopy of adult CX3CR1+/gfp mice (n = 50–60 cells per group of four mice in three independent experiments). GFP+ F4/80+ Macs co-localizing with β 3-tubulin+ nerves, GFP+ F4/80+ interstitial Macs (partly Ly6C +), and other interstitial F4/80+ cells were analyzed for GFP fluorescence intensity. Dotted lines indicate thresholds for qualitative subset discrimination in microscopy. One-way ANOVA, Dunnett correction, $p < 0.01$.

(E) IF staining and mean fluorescence intensity (MFI) in flow cytometry for CD206. Surrounding CD206+ cells were F4/80+ (see Figure S4G). Scale bars: 50 μ m.

(F) 3D reconstruction of CX3CR1hi Macs and a β 3-tubulin + nerve. Scale bars: 10 μ m. Volume: 30 μ m.

(G) Nerve with associated Macs (white arrows). The maximum intensity projection of 110 μ m tissue volume includes CX3CR1+ $\gamma\delta$ T cells (green triangle, exemplary) and hair follicles (white triangle, exemplary) of the epidermis. h, hair; black circle, air bubble. Scale bar: 500 μ m.


(H) Density of CX3CR1hi Macs on nerves (in Macs/mm) correlated to nerve bundle diameter (n = 4 mice).

(I) Distance between single CX3CR1hi Macs depending on nerve bundle diameter (n = 4 mice).

(J) CX3CR1hi Macs co-localized to sensory nerves (CGRP+). Scale bars: 25 μ m.

Data are mean $\pm$ SEM. $p < 0.5$, $p < 0.01$ (two-tailed unpaired t test). Data are representative of at least three mice in two or more independent experiments.


Fig 4. PMID: 31201094



Immunofluorescence Microscopy

P2X7-Rs are highly expressed in ON-RGCs of Thy1-YFP-H mice. Thy1-YFP-H RGCs (green) are projections of the original confocal Z-stacks, while the P2X7-R puncta contacting the dendrites (red) are digitally rendered as spots (presented as 3 times larger than the true colocalised puncta for ease of viewing). (A–C) Thy1-YFP-H labelled ON-RGC (green) showing dense labelling for P2X7-R puncta (red) shown in (A) orthogonal view, (B) from above and (C) magnified inset from (B). (D) A single plane of raw data from the ON-cell (green) with P2X7-R staining (red). (E–G) Thy1-YFP-H labelled OFF-RGC (green) showing sparser labelling for P2X7-R puncta (red) than ON-RGCs. The OFF-RGC is shown in (E) orthogonal view, (F) from above and (G) magnified inset from F. (H–L) Thy1-YFP-H labelled ON–OFF RGC (green) in the orthogonal view (H) displaying dendritic fields stratifying in ON and OFF sublaminae. Labelling for the P2X7-R puncta (red) in the ON sublamina (I) from above and (J) magnified inset from (I). Labelling for the P2X7-R puncta (red) in the OFF sublamina (K) from above and (L) magnified inset from K. (M) Quantification of P2X7-R puncta in the ON and OFF lamina of the IPL for ON-, OFF- and ON–OFF RGCs. Numbers indicate number of cells from 4 animals. * $p < 0.01$; one-way ANOVA, Tukey's multiple comparisons test. Scale bar is 10 μm.

Fig 6. PMID: 33603067




Immunofluorescence Microscopy

(C) Passive diffusion from the cytoplasm to the nucleus measured by FRAP, in HeLa cells transduced with mTagBFP-2A control, SUN1 or SUN2 lentivectors and a lentivector encoding GFP. Top, representative layout of a photobleached control HeLa cell: imaging starts at $t = 0$ after high intensity laser exposure and ends at $t = 80$ seconds, when fluorescence in the nucleus has been recovered (scale bar: 5 μ m). Bottom left: GFP intensity recovery over time (s). For each individual cell, the background intensity was subtracted and intensity was normalized to 0 at t_0 after photobleaching while the max intensity reached during the course of each measurement was set to 1. Curves show mean and standard deviation per time point across indicated number of cells per condition. Bottom right, hillslopes for each cell were calculated via non-linear regression fit (total number of nuclei analyzed per cell line are indicated, $n=1$ experiment; un-paired ANOVA one-way with Dunnett's post-test, line at mean $\pm$ SEM). FS5. PMID: 34592156



Immunofluorescence Microscopy

Multi-color Immunofluorescence Microscopy using DyLight™ dyes. Tissue: human breast carcinoma. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: Anti-Histone and Anti-Tubulin antibody at 10 μ g/mL for 1 h at RT. Secondary antibody: DyLight™ 488 conjugate and DyLight™ 549 conjugate goat secondary antibody at 1:10,000 for 45 min at RT. Localization: Histone is nuclear and Tubulin is cytoplasmic. Staining: Anti-Histone detection using a DyLight™ 488 conjugate (green) fluorescent signal and Anti-Tubulin was detected using a DyLight™ 549 conjugate (red) fluorescent signal. Nuclei were counter-stained using DAPI (blue).




Diagram

Comparison of fluorescence absorption and emission spectra for DyLight™ 488 and Alexa Fluor 488 in PBS, pH7.2. The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation.

Diagram

Properties of DyLight™ Fluorescent Dyes. The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation.

Emission	Color	DyLight™ Dye	Ex/Em (nm)	ϵ (M ⁻¹ cm ⁻¹)	Similar Dyes
Blue		405	400/420	30,000	Alexa™ 405, Cascade Blue
Green		488	493/518	70,000	Alexa™ 488, Cy2®, FITC
Yellow		549	550/568	150,000	Alexa™ 546, Alexa 555, Cy3®, TRITC
Red		649	646/674	250,000	Alexa™ 647, Cy5®
Near Infrared		680	682/715	140,000	Alexa™ 680, Cy5.5®, IRDye™ 700
Infrared		800	770/794	270,000	IRDye™ 800



Western Blot

Western Blot of DyLight™ 488 conjugated anti-GFP antibody to detect GFP control proteins. Lane 1: His-Sumo-GFP. Lane: Molecular Weight. Lane 2: Beta-Galactosidase (negative control). Lane 3: recombinant GFP control protein (000-001-215). Load: 35 µg per lane. Primary antibody: none. Secondary antibody: DyLight™ 488 conjugated anti-GFP goat secondary antibody at 1:5,000. Block: MB-070 for 2 hr at RT. Predicted/Observed size: 27kDa/54kDa, 27kDa for rGFP/~45kDa His-Sumo-GFP. Other band(s): none.

References

- Ginn SL et al. Functional editing of the OTC locus by targeted integration with phenotype correction and restoration of endogenous expression patterns. *Mol Ther.* (2026)
- Świst-Rosowska KM et al. The basic domain of Suv39h2 buffers mitoxantrone-induced heterochromatin destabilization. *iScience.* (2026)
- Kolter J et al. Sensory neurons shape local macrophage identity via TGF- β signaling. *Immunity.* (2025)
- Park J et al. Generation and characterization of a zebrafish knockout model of abcb4, a homolog of the human multidrug efflux transporter P-glycoprotein. *Hum Genomics.* (2023)
- Kim H et al. Reactive astrocytes transduce inflammation in a blood-brain barrier model through a TNF-STAT3 signaling axis and secretion of alpha 1-antichymotrypsin. *Nat Commun.* (2022)
- Annunziata F et al. Paneth cells drive intestinal stem cell competition and clonality in aging and calorie restriction. *Eur J Cell Biol.* (2022)
- Lee, PY et al. The Effect of Aging on Retinal Function and Retinal Ganglion Cell Morphology Following Intraocular Pressure Elevation. *Frontiers in Aging Neuroscience* (2022)
- Paterson, N et al. Macrophage network dynamics depend on haptokinesis for optimal local surveillance. *ELife* (2022)
- Kim H et al. Generation of human pluripotent stem cell-derived fused organoids with oligodendroglia and myelin. *STAR Protoc.* (2021)
- Freter R et al. Establishment of a fluorescent reporter of RNA-polymerase II activity to identify dormant cells. *Nat Commun.* (2021)
- Bhargava A et al. Inhibition of HIV infection by structural proteins of the inner nuclear membrane is associated with reduced chromatin dynamics. *Cell Rep.* (2021)
- Paterson N et al. Macrophage network dynamics depend on haptokinesis for optimal local surveillance. *bioRxiv Preprint* (2021)
- Wang AYM et al. Retinal ganglion cell dysfunction in mice following acute intraocular pressure is exacerbated by P2X7 receptor knockout. *Sci Rep.* (2021)
- Xu R et al. OLIG2 drives abnormal neurodevelopmental phenotypes in human iPSC-based organoid and chimeric mouse models of down syndrome. *Cell Stem Cell.* (2019)
- Gentili et al. The N-Terminal Domain of cGAS Determines Preferential Association with Centromeric DNA and Innate Immune Activation in the Nucleus. *Cell Reports* (2019)
- Kolter J et al. A subset of skin macrophages contributes to the surveillance and regeneration of local nerves. *Immunity* (2019)
- Begolly et al. Fractionation enhances acute oligodendrocyte progenitor cell radiation sensitivity and leads to long term depletion. *Glia* (2018)
- Haas et al. Viral targeting of TFIIIB impairs de novo polymerase II recruitment and affects antiviral immunity. *PLOS Pathogens* (2018)
- Fletcher EL et al. Ganglion Cell Assessment in Rodents with Retinal Degeneration. *Methods Mol Biol.* (2018)
- Russo MV et al. Distinct myeloid cell subsets promote meningeal remodeling and vascular repair after mild traumatic brain injury. *Nat Immunol.* (2018)

- Bongianino R et al. Allele-Specific Silencing of Mutant mRNA Rescues Ultrastructural and Arrhythmic Phenotype in Mice Carriers of the R4496C Mutation in the Ryanodine Receptor Gene (RYR2). *Circ Res.* (2017)
- Dufour et al. Direct Correlation between Motile Behavior and Protein Abundance in Single Cells. *PLOS Computational Biology* (2016)
- Dufour YS et al. Direct correlation between motile behavior and protein abundance in single cells. *PLoS Comput Biol.* (2016)
- Ho T et al. Immunolocalization of the P2X4 receptor on neurons and glia in the mammalian retina. *Neuroscience.* (2014)
- Chang THT et al. New insights into lineage restriction of mammary gland epithelium using parity-identified mammary epithelial cells. *Breast Cancer Res.* (2014)
- Liu Y et al. OLIG gene targeting in human pluripotent stem cells for motor neuron and oligodendrocyte differentiation. *Nat Protoc.* (2011)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.