

Datasheet for 600-401-378

V5 Epitope Tag Antibody

Overview

Description:	Anti-V5 Epitope Tag (RABBIT) Antibody - 600-401-378
Item No.:	600-401-378
Size:	100 µg
Applications:	ELISA, WB, IP
Reactivity:	V5-Tag
Host Species:	Rabbit

Product Details

Background: Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells. Rockland Immunochemicals produces anti-epitope tag antibodies against many common epitope tags including Myc, GST, GFP, 6X His, MBP, FLAG, HA and V5. Rockland Immunochemicals also produces antibodies to other tags including FITC, Rhodamine (TRITC), DNP and biotin.

Synonyms:	Rabbit Anti-V5 Epitope Tag Antibody, Rabbit Anti V5 Epitope Tag Antibody, Rabbit Anti-V5 Tag
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	V5-Tag
--------------------	--------

Immunogen Type:	Conjugated Peptide
Immunogen:	This antibody was purified from whole rabbit serum prepared by repeated immunizations with V5 epitope tag peptide corresponding to aa 95-108 of the V protein conjugated to KLH using maleimide.
Purity/Specificity:	This affinity-purified antibody is directed against V5 motif and is useful in determining its presence in various assays. This polyclonal anti-V5-tag antibody detects over-expressed proteins containing the V5 epitope tag. To date this antibody has reacted with all V5 tagged proteins tested so far. In western blotting of bacterial extracts the antibody does not cross-react with endogenous proteins. The antibody recognizes the V5-epitope tag (GKPIPPLLGLDST) fused to either the carboxy- terminal end of targeted proteins in transfected or transformed cells. Although not yet tested, expect reactivity with recombinant proteins prepared with the V5-epitope tag fused to the amino terminal end as well.
Relevant Links:	<ul style="list-style-type: none">• NCBI - 55775699

Application Details

Tested Applications:	ELISA, WB
Suggested Applications:	IP (Based on references)
Application Note:	Anti-V5 is optimally suited for monitoring expression of V5-tagged fusion proteins. The V5 epitope tag is derived from a small epitope (Pk) present on the P and V proteins of the paramyxovirus of simian virus 5 (SV5). The V5 tag is usually used with all 14 amino acids (GKPIPPLLGLDST), although it has also been used with a shorter 9 amino acid sequence (IPNPLLGLD). This antibody has been tested by ELISA and western blotting against both the immunizing peptide and V5 containing recombinant proteins. Although not tested, this antibody is likely functional for immunoprecipitation and immunocytochemistry.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:60,000
IHC:	1:500 - 1:3,000
WB:	1:5,000 - 1:10,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.8 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images




Western Blot

Autoprocessing of precursors with different C-terminal tags. All fusion precursors were derived from GST-M2-PRNL-Flag for specific examination of P site cleavage reaction. (A) The sequences at the junction between the PR and the C-terminal tag of some constructs are listed. (B) Autoprocessing of precursors carrying different small peptide epitopes or no tag were analyzed using a mouse epitope-specific antibody (Flag, Myc, HA), polyclonal rabbit anti-V5, or anti-PR as underlined at left, and GST primary antibodies for dual visualization. Because the signal detected by anti-PR antibody was weak, the boxed region was enhanced to show the mature protease (bottom). Autoprocessing of precursors fused to a GCN4 motif (C) or to reverse transcriptase (RT) in which the native cleavage site between PR and RT remained cleavable (E) or was mutated to uncleavable (D) were also compared. Asterisks indicate self degradation IC50 of the PR-containing product; open triangles indicate the apparent IC50 of the cleavage reaction. The expected products from P site cleavage reaction are connected by a solid line. Fig 5. PMID: 21985091




Western Blot

B, Western analysis with anti-V5 tag monoclonal antibody showing exclusive expression of V5-tagged mutant or WT KREPB5 proteins from the β -tubulin locus in BF and PF CN cells following repression (R) of the tet-regulatable WT KREPB5 allele for 3 or 4 days, respectively (2×10^6 cells/lane). TAP-tagged WT KREPB5 is detected with PAP antibody in PF cells in which the tet-regulatable allele is expressed (E). Fig 2. PMID: 26304125




Immunoprecipitation

Effects of KREPB5 PUF motif mutations on BF and PF editosomes. A) anti-V5 immunoprecipitation (IP) of exclusively expressed V5-tagged mutant or WT KREPB5 from PF and BF cells in which the tet-regulatable WT KREPB5 allele was repressed for 4 (PF) or 3 (BF) days. Left panel, anti-V5 immunoprecipitates from WT or mutant KREPB5 from PF and BF (6.25 and 12.5% of each immunoprecipitate, respectively) probed with monoclonal antibodies against KREPA1, KREPA2, KREL1, and KREPA3 (A1, A2, L1, and A3) and with anti-V5 tag antibody (V5). No antibody (Mock) and untagged CN cell lysates are controls. Middle and right panels, comparisons of 6.25% of the PF immunoprecipitates and PF controls with 2.5% of cell lysate samples from which the immunoprecipitates were derived. Fig 5. PMID: 26304125



Western Blot

(A) HCT116 β -catenin^{WT}Clover/ Δ 45Cherry (clone #37) cells express comparable amounts of β -catenin to the parental HCT116 WT cells. (B–D) Immunoprecipitation using HCT116 β -catenin^{WT}Clover/ Δ 45 (clone #33 – left), β -catenin^{WT}/ Δ 45Cherry (clone #45 – middle), and β -catenin^{WT}Clover/ Δ 45 (clone #24 – right) were performed with GFP/Cherry or control beads, followed by Western blotting with the indicated antibodies. Representative results from three independent experiments are shown. SF2. PMID: 26304125



Western Blot


(C) BF cells that exclusively expressed V5-tagged WT or mutant B4, or untagged WT B4 control following two days of repression of the regulatable WT B4 allele were used for anti-V5 tag immunoprecipitation of B4-bound protein complexes. Cleared input cell lysates (5%) and anti-V5 immunoprecipitates (10%) were analyzed by Western blotting. Blots were probed with monoclonal antibodies against RECC proteins KREPA1, KREPA2, KREL1 and KREPA3, and anti-V5 antibody. Fig 6. PMID: 38650737

Western Blot

Wnt ligand production increases Evi protein levels. Wild-type (wt) or EviKO HEK293T cells were transfected with Wnt3A or IGFBP5-V5 expression plasmids and subjected to Western blot analysis. Specific Evi bands are indicated by arrows and unspecific bands by asterisks. Endogenous Evi is not only detectable as a monomeric form (46 kDa) but also as SDS-resistant dimers (80 kDa). Clonal EviKO HEK293T cells were generated via CRISPR/Cas9 using Evi sgRNA #2 (EviKO2.9) or Evi sgRNA #1 (EviKO1.1; Appendix Fig S2). Figure 1B. Source: EMBO J, PMID: 29378775.


Western Blot

Evi stabilization is dependent on Wnt palmitoylation. Western blot analysis of endogenous Evi in wt, PorcnKO, or EviKO HEK293T cells upon overexpression of Wnt3A or IGFBP5-V5. PorcnKO1.2 and PorcnKO1.4 indicate clone #2 and clone #4 of PorcnKO HEK293T cells generated with Porcn sgRNA1 (Appendix Fig S3). Clonal EviKO HEK293T cells were generated with Evi sgRNA2 (EviKO2.9; clone #9) or Evi sgRNA1 (EviKO1.1; clone #1; Appendix Fig S2). Increase in total β -catenin protein served as control for Wnt pathway activation. Figure 2C. Source: EMBO J, PMID: 29378775.




Western Blot

Evi stabilization is dependent on Wnt palmitoylation. Western blot analysis of endogenous Evi in HEK293T cells transfected with the indicated overexpression plasmids. When indicated, the cells were additionally treated with 5 μ M LGK974 for 48 h. Figure 2D. Source: EMBO J, PMID: 29378775.





Western Blot

Evi is ubiquitinated and degraded from the ER by the proteasome. HEK293T cells were transfected with Wnt3A or IGFBP5-V5 expression plasmids and treated for 24 h with DMSO, MG132, or bortezomib in the indicated concentrations. Figure EV3B. Source: EMBO J, PMID: 29378775.



Western Blot

Evi poly-ubiquitination is regulated by the presence of Wnt proteins. HEK293T cells were transfected with the indicated plasmids and additionally treated with 5 μ M LGK974 for 48 h when indicated. In case of Wnt3A-KDEL, the ER-retaining sequence KDEL was C-terminally fused to Wnt3A. Dvl2-HA, Dvl3, and β -catenin-YFP overexpression was used as negative control to verify that Evi stabilization was not due to downstream activation of Wnt signaling. All Western blots are representative of three independent experiments. β -Actin was used as loading control. Specific Evi bands are marked by arrows and unspecific bands by asterisks. Figure 3D. Source: EMBO J, PMID: 29378775.



Western Blot

Evi is not transcriptionally regulated by Wnt. D'Twenty-four hours after reverse transfection with Ctrl or CTNNB1 siRNA, HEK293T cells were transfected with Wnt3A or IGFBP5-V5 expression plasmids and analyzed (D) for the indicated proteins via immunoblotting or (D') for canonical Wnt activity using the TCF-Luciferase Wnt reporter assay. Immunoblotting is representative of three independent experiments, and Wnt reporter activity was calculated as mean from three independent experiments \pm s.d. Figure EV1D. Source: EMBO J, PMID: 29378775.

Western Blot

Anti-V5 epitope tag polyclonal antibody detects V5-tagged recombinant protein by western blot. This antibody was used at 1.0 μ g/ml to detect 0.05 μ g (lane 2) of full-length recombinant mouse serum albumin containing the V5 epitope tag at the carboxy end. Comparison to MW markers (lane 1) indicates detection of monomeric V5 tagged albumin. A 4-20% gradient gel was used to separate the protein by SDS-PAGE under non-reducing conditions. The protein was transferred to nitrocellulose using standard methods. After blocking the membrane was probed with the primary antibody overnight at 4 $^{\circ}$ C followed by washes and reaction with a 1:10,000 dilution of IRDye 800 conjugated Gt-a-Rabbit IgG [H&L] (code 611-132-122) for 45 min at room temperature. LICOR's Odyssey[®] Infrared Imaging System was used to scan and process the image. Other detection systems will yield similar results.

References

- McDermott SM et al. Deep mutational scanning of the RNase III-like domain in *Trypanosoma brucei* RNA editing protein KREP4. *Front Cell Infect Microbiol.* (2024)
- Ambrosi G et al. Allele-specific endogenous tagging and quantitative analysis of β -catenin in colorectal cancer cells. *Elife.* (2022)
- Kotredes KP et al. Characterization of cancer-associated IDH2 mutations that differ in tumorigenicity, chemosensitivity and 2-hydroxyglutarate production. *Oncotarget.* (2019)
- Glaeser et al. ERAD-dependent control of the Wnt secretory factor Evi. *The EMBO Journal* (2018)
- Watanabe et al. The HIV-1 late domain-2 S40A polymorphism in antiretroviral (or ART)-exposed individuals influences protease inhibitor susceptibility. *Retrovirology* (2016)
- McDermott SM et al. Differential editosome protein function between life cycle stages of *Trypanosoma brucei*. *J Biol Chem.* (2015)
- Huang L et al. Flexible catalytic site conformations implicated in modulation of HIV-1 protease autoprocessing reactions. *Retrovirology.* (2011)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.