

Datasheet for 600-401-458**BACH1 Antibody****Overview**

Description:	Anti-BACH1 (RABBIT) Antibody - 600-401-458
Item No.:	600-401-458
Size:	100 µg
Applications:	ELISA, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	BACH1 (also known as BRCA1 interacting protein C-terminal helicase 1, BRCA1-interacting protein 1 and BRCA1-associated C-terminal helicase 1) is a member of the RecQ DEAH helicase family and interacts with the BRCT repeats of breast cancer, type 1 (BRCA1). The bound complex is important in the normal double-strand break repair function of breast cancer, type 1 (BRCA1). The BACH1 gene may be a target of germline cancer-inducing mutations. BACH1 is localized within the nucleus and functions as a DNA-dependent ATPase and 5' to 3' DNA helicase. Two isoforms have been identified for this protein.
Synonyms:	rabbit anti-BACH1 antibody, BACH-1, BACH 1, Fanconi anemia group J protein, FACJ, BRAC 1 Associated C Terminal Helicase 1 antibody, BRCA 1 Interacting Protein 1 antibody, BRCA1 binding helicase like protein BACH1 antibody, BRCA1 interacting protein C terminal helicase 1 antibody, BRIP 1 antibody, ATP-dependent RNA helicase BRIP1, FANCI, FANCI-J, FANCI J
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	BRIP1
Reactivity:	Human
Immunogen Type:	Recombinant Protein

Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a recombinant protein corresponding to amino acids 92-104 of isoform 1 of human BACH1 protein.
Purity/Specificity:	This affinity purified antibody is directed against human BACH1 protein. The product was affinity purified from monospecific antiserum by immunoaffinity purification. A BLAST analysis was used to suggest cross reactivity with BACH1 protein from human (100% homology) and chimpanzee (92% homology). Expect reactivity with isoform 1 and isoform 2 of BACH1. Reactivity against BACH1 homologues from rat and mouse is not expected. Reactivity against homologues from other sources is not known.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - Q9BX63• NCBI - 14042978• GenelD - 83990

Application Details

Tested Applications:	ELISA, WB
Application Note:	This affinity purified antibody has been tested for use in ELISA and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 105 - 140 kDa in size corresponding to isoforms of BACH1 protein by western blotting in the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:44,000
WB:	1:500- 1:2,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western blot using Rockland's Affinity Purified anti-BACH1 antibody shows detection of a band at ~105 kDa (lane 1) corresponding to human BACH1 present in a 293 whole cell lysate (p/n W09-000-365). Approximately 35 µg of lysate was separated on a 4-20% Tris-Glycine gel by SDS-PAGE and transferred onto nitrocellulose. After blocking the membrane was probed with the primary antibody diluted to 1:1,000. Reaction occurred 2 h at room temperature followed by washes and reaction with a 1:10,000 dilution of IRDye™800 conjugated Gt-a-Rabbit IgG [H&L] MX (p/n 611-132-122) for 45 min at room temperature (800 nm channel, green). Molecular weight estimation was made by comparison to prestained MW markers in lane M (700 nm channel, red). IRDye™800 fluorescence image was captured using the Odyssey® Infrared Imaging System developed by LI-COR. IRDye is a trademark of LI-COR, Inc. Other detection systems will yield similar results.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.