

Datasheet for 600-401-853**GGA1 Antibody****Overview**

Description:	Anti-GGA1 (RABBIT) Antibody - 600-401-853
Item No.:	600-401-853
Size:	100 µg
Applications:	ELISA, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	GGA1 (also known as ADP-ribosylation factor binding protein, Golgi-localized, gamma ear-containing, ARF-binding protein 1, Gamma-adaptin related protein 1) plays a role in protein sorting and trafficking between the trans-Golgi network (TGN) and endosomes. This protein mediates the ARF-dependent recruitment of clathrin to the TGN and binds ubiquitinated proteins and membrane cargo molecules with a cytosolic acidic cluster-dileucine (AC-LL) motif. GGA1 is a monomeric protein that interacts with NECAP1, GGA2 and GGA3 as well as RABEP1, RABGEF1 and type-I membrane proteins SORT1, SORL1, LRP3, M6PR/CD-MPR, IGF2R/CI-MPR and BACE1. GGA1 is a ubiquitously membrane-associated protein that is localized to TGN and endosomes. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Synonyms:	rabbit anti-GGA-1 rabbit, GGA 1,GGA1, ADP-ribosylation factor binding protein, Golgi-localized gamma ear-containing ARF-binding protein 1, Gamma-adaptin related protein 1
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	GGA1
Reactivity:	Human
Immunogen Type:	Conjugated Peptide

Immunogen:	This affinity-purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near amino acids 330-360 of Human GGA-1.
Purity/Specificity:	This affinity purified antibody is directed against human GGA1 protein. The product was affinity purified from monospecific antiserum by immunoaffinity purification. A BLAST analysis was used to suggest reactivity with this protein from human based on 100% homology for the immunogen sequence. Cross reactivity may occur with GGA1 from bovine (87% homology), mouse (82% homology) and rat (76% homology) sources. The immunizing sequence is present only in isoform 1 of this protein, therefore cross reactivity with splice variants is not expected. Cross reactivity with GGA1 homologues from other sources has not been determined.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - Q9UJY5• NCBI - 9558729• GenelD - 26088

Application Details

Tested Applications:	ELISA, WB
Application Note:	This affinity purified antibody has been tested for use in ELISA and western blot. Specific conditions for reactivity should be optimized by the end user. Expect bands at 70 kDa in size corresponding to GGA1 by western blotting in the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:50,000
IP:	1:100
WB:	1:500 - 1:3,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.35 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western blot using Rockland's Affinity Purified anti-GGA1 antibody shows detection of bands at ~100 kDa corresponding to YFP-GGA1 fusion present in a lysate of HEK293 cells over-expressing the recombinant protein (arrowhead). Approximately 35 µg of lysate was separated on a 4-20% gel by SDS-PAGE and transferred onto nitrocellulose. After blocking the membrane was probed with the primary antibody diluted to 1:1,350. Reaction occurred overnight at 4° followed by washes and reaction with a 1:10,000 dilution of IRDye™800 conjugated Gt anti-Rb for 45 min at room temperature. IRDye™800 fluorescence image was captured using the Odyssey® Infrared Imaging System developed by LI-COR. IRDye is a trademark of LI-COR, Inc. Other detection systems will yield similar results.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.