

Datasheet for 600-401-876**APC6 phospho T580 Antibody****Overview**

Description:	Anti-APC6 pT580 (RABBIT) Antibody - 600-401-876
Item No.:	600-401-876
Size:	100 µg
Applications:	ELISA, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	APC6 (also known as Cell division cycle protein 16 homolog (CDC16Hs), anaphase promoting complex subunit 6, and cyclosome subunit 6) is a component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis and the G ₁ phase of the cell cycle. The APC/C is composed of at least 11 subunits. APC6 interacts with PPP5C and CDC20. This protein colocalizes with CDC27 to the centrosome at all stages of the cell cycle and to the mitotic spindle. Splice variants have been described for this protein.
Synonyms:	rabbit anti-APC6 pT580 Antibody, APC 6, APC-6, Anaphase promoting complex subunit 6 antibody, CDC16Hs antibody, Cell division cycle 16 homolog antibody, Cell division cycle protein 16 homolog antibody, Cyclosome subunit 6 antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	CDC16
Reactivity:	Human
PTM Specificity:	Phosphorylation
Immunogen Type:	Conjugated Peptide

Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region surrounding pT580 of Human Apc6 protein.
Purity/Specificity:	This product is an affinity purified antibody produced by immunoaffinity chromatography using phospho peptide coupled to agarose beads followed by solid phase adsorption(s) against non-phospho peptide and non-specific peptide to remove any unwanted reactivities. This antibody is specific for phosphorylated human APC6 protein at the pT580 residue. BLAST analysis indicates 100 % homology of the immunizing sequence with APC6 from human. No cross reactivity is expected against homologues from other sources. This antibody is expected to react with all isoforms of this protein. Minimal reactivity is expected with the non-phosphorylated form of the protein.
Relevant Links:	• NCBI - 37537763 • UniProtKB - Q13042 • GeneID - 8881

Application Details

Tested Applications:	ELISA, WB
Application Note:	This affinity purified antibody has been tested for use in ELISA and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band ~ 72 kDa, 71.5 kDa or 65.9 kDa in size corresponding to isoforms of APC6 by western blotting in the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:1,000 - 1:10,000
IF:	User Optimized
WB:	1:500 - 1:3,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.97 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western blot using Rockland's Affinity Purified anti-APC6 pT580 antibody shows detection of a band ~72 kDa corresponding to phosphorylated human APC6 (arrowhead lane 1). Lane 1 - nocodazole treated HeLa whole cell lysate (p/n W09-001-A81). Lane 2 - Reactivity is not seen in lysates from asynchronous HeLa whole cell cultures (p/n W09-000-364). Each lane contains approximately 35ug of lysates, separated by 4-20% SDS-PAGE Tris-HEPES and then transferred to nitrocellulose. After blocking the membrane was probed with the primary antibody diluted to 1:1,000 overnight at 4°C followed by washes and reaction with a 1:10,000 dilution of IRDye™800 conjugated Gt-a-Rabbit IgG [H&L] MX (p/n 611-132-122) for 45 min at room temperature. IRDye™800 fluorescence image was captured using the Odyssey® Infrared Imaging System developed by LI-COR. IRDye is a trademark of LI-COR, Inc. Other detection systems will yield similar results.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.