

Datasheet for 600-401-989**c-Met phospho Y1349 / pY1356 Antibody****Overview**

Description:	Anti-c-MET pY1349 pY1356 (RABBIT) Antibody - 600-401-989
Item No.:	600-401-989
Size:	100 µg
Applications:	ELISA, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	This antibody is designed, produced, and validated as part of a collaboration between Rockland and the National Cancer Institute (NCI) and is suitable for Cancer, Immunology and Nuclear Signaling research. Anti-c-MET is the receptor for hepatocyte growth factor (also known as scatter factor, HGF/SF), and belongs to the tyrosine kinase superfamily. Interaction of c-Met with HGF results in autophosphorylation of c-Met at multiple tyrosines. Phosphorylation of Y1234/1235 in the c-Met kinase domain is critical to kinase activation. When phosphorylated, Y1349 and Y1356, along with surrounding amino acids, form a unique bidentate docking site for substrates such as Gab1, Grb2, phosphatidylinositol 3-kinase (PI3K) and others. C-Met mainly uses the Gab1 scaffolding adaptor in its initial step of signal transmission. Well-characterized downstream signaling pathways that are activated by c-Met include the ERK/MAPK, PI3K-Akt/PKB, Crk-Rap and Rac-Pak pathways, resulting in proliferation and increased cell survival. Anti-Phospho cMET was developed with NCI and is ideal for Cancer Research.
Synonyms:	rabbit anti-c-Met pY1349/pY1356 Antibody, rabbit anti-c-Met dual phosphorylated Antibody, HGF receptor antibody, HGF/SF receptor antibody, HGFR antibody, MET antibody, Met proto oncogene tyrosine kinase antibody, SF receptor, Tyrosine-protein kinase Met, Scatter factor receptor, Proto-oncogene c-Met, Hepatocyte growth factor receptor
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	MET
-------------------	-----

Reactivity:	Human
PTM Specificity:	Dual Modification
Immunogen Type:	Conjugated Peptide
Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic phospho-peptide corresponding to residues surrounding Y1349 and Y1356 of human c-Met protein.
Purity/Specificity:	Phospho c-MET Antibody was produced from monospecific antiserum by immunoaffinity chromatography using dual-phospho-peptide coupled to agarose beads followed by cross-adsorption against nonphospho-peptide. Antibody is specific for human c-Met protein phosphorylated at Y1349 and Y1356. A BLAST analysis was used to suggest cross-reactivity with c-Met from human, mouse and rat based on 100% homology with the immunizing sequence. Cross-reactivity with c-Met from other sources has not been determined.
Relevant Links:	• NCBI - 42741655 • UniProtKB - P08581 • GenelD - 4233

Application Details

Tested Applications:	ELISA, WB
Application Note:	This affinity purified antibody has been tested for use in ELISA and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 150 kDa in size corresponding to phosphorylated c-Met protein by western blotting in the appropriate cell lysate or extract. This phospho-specific polyclonal antibody reacts with human c-Met pY1349pY1356 and shows minimal reactivity by ELISA against the non-phosphorylated form of the immunizing peptide.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:93,000
WB:	1µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.96 mg/ml
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store Phospho antibody at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

B) MM cells were treated or not with 50μM TDYA for 36 h and probed in immunoblotting with the indicated antibodies. Immunoblots demonstrate representative images of at least two independent experiments with no substantial experimental variability. Fig 5. PMID: 39885295




Western Blot

C) MM cells were treated or not with 50μM TDYA for 36 h and probed in immunoblotting with the indicated antibodies. Immunoblots demonstrate representative images of at least two independent experiments with no substantial experimental variability. Fig 5. PMID: 39885295



Western Blot

Western blot using Rockland's affinity purified anti-c-Met pY1349pY1356 antibody shows detection of phosphorylated c-Met. Human mammary B5/589 epithelial cells were serum-deprived and treated with or without HGF. Cell lysates were immunoprecipitated with the anti-c-Met antibody, resolved by SDS-PAGE, transferred to PVDF membrane, and probed with anti-c-Met pY1349pY1356. Personal communication, D. Bottaro and T. Ito, NCI, Bethesda, MD



Western Blot

Western blot using Rockland's affinity purified anti-c-Met pY1349pY1356 antibody shows detection of phosphorylated c-Met. Human mammary epithelial cells (B5/589) were serum deprived and stimulated with (+) and without (-) HGF. Cell lysates were immunoprecipitated using a human anti-c-Met antibody, resolved by SDS-PAGE, transferred to PVDF membrane and probed using various antibodies. Lane 1 and 2 where probed using the anti-c-Met pY1349pY1356 antibody (#600-401-989), lane 3 and 4 where probed using an anti-phosphotyrosine antibody as phosphorylation control and lane 5 and 6 where probed using an anti-c-Met antibody as a total Met loading control. Bands recognized at an apparent molecular weight of ~170 kDa are total Met, ~145 kDa phosphorylated Met beta chain in HGF +, ~150 kDa phosphorylated Met in +/- HGF, ~50kDa phosphorylated Met alpha chain. Personal communication with D. Bottaro and F. Cecchi, CCR-NCI, Bethesda, MD.

References

- Han Z et al. Targeting ABCD1-ACOX1-MET/IGF1R axis suppresses multiple myeloma. *Leukemia*. (2025)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.