

Datasheet for 600-401-A37**Wnt1 Antibody****Overview**

Description:	Anti-Wnt1 (RABBIT) Antibody - 600-401-A37
Item No.:	600-401-A37
Size:	100 µg
Applications:	ELISA, WB
Reactivity:	Human, Mouse
Host Species:	Rabbit

Product Details

Background:	The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. Wnt1 (Wingless-type MMTV integration site family member 1) is a member of the WNT gene family. It is highly conserved in evolution and the protein encoded by this gene is known to be 98% identical to mouse Wnt1 protein at the amino acid level. Studies in mouse indicate that the Wnt1 protein functions in the induction of the mesencephalon and cerebellum. This gene was originally considered as a candidate gene for Joubert syndrome, an autosomal recessive disorder with cerebellar hypoplasia as a leading feature. However, further studies suggested that the gene mutations might not have a significant role in Joubert syndrome. Wnt1 is secreted as an extracellular matrix protein.
Synonyms:	rabbit anti-WNT-1 antibody, rabbit anti-WNT1 antibody, INT1 antibody, Murine mammary tumor virus integration site 1 antibody, Oncogene INT1 antibody, Proto oncogene protein Wnt 1 antibody, Wingless type MMTV integration site family member 1 antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	WNT1
Reactivity:	Human, Mouse

Immunogen Type:	Conjugated Peptide
Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region of human Wnt1 protein.
Purity/Specificity:	This product was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with human and mouse Wnt1 protein. A BLAST analysis was used to suggest cross-reactivity with Wnt1 from mouse, human, rat, bovine, dog, macaque, and opossum based on a 100% homology with the immunizing sequence. Partial cross-reactivity is expected against chicken Wnt1 based on a 91% sequence homology. Cross-reactivity with Wnt1 from other sources has not been determined.
Relevant Links:	• UniProtKB - P04628 • NCBI - 4885655 • GeneID - 7471

Application Details

Tested Applications:	ELISA, WB
Application Note:	This affinity purified antibody has been tested for use in ELISA and western blotting.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:25,000
WB:	1:1,500 - 1:6,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None




Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Expression of Wnt1 in kidney homogenates of sham and nephrectomized rats. The expression of Wnt1 in kidney homogenates of sham and nephrectomized rats 18 weeks after the surgery was analyzed by immunoblotting. Blotting of GDPH served as loading control. Results are derived from 4-5 animals per group. Figure provided by CiteAb. Source: Biomed Res Int, PMID: 24995284.

Western Blot

Western blot using Rockland's affinity purified anti-Wnt1 antibody shows detection of endogenous Wnt1. Lane 1: human-derived MCF7 cell lysate (p/n W09-000-360). Lane 2: mouse-derived 3T3 cell lysate (p/n W10-000-358). The band at ~41kDa, indicated by the arrowhead, corresponds to Wnt1. After transfer, the membrane was blocked with 5% BLOTTO (p/n B501-0500). Primary antibody was used at a 1:1,400 dilution in PBS containing 1% BLOTTO. The specificity of the antibody was confirmed by peptide competition which completely blocked reaction of the antibody with Wnt1 (data not shown).

References

- Banon-Maneus E et al. Wnt pathway activation in long term remnant rat model. *Biomed Res Int.* (2014)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.