

Datasheet for 600-401-AB9**CARMA2 Antibody****Overview**

Description:	Anti-CARMA2 (RABBIT) Antibody - 600-401-AB9
Item No.:	600-401-AB9
Size:	100 µg
Applications:	ELISA, IHC
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	CARMA proteins belong to the membrane-associated guan-ylate kinase-like (MAGUK) family of proteins that can function as molecular scaffolds that assist assembly of signal transduction molecules. CARMA1, CARMA2, and CARMA3 share high degrees of sequence and functional homology, but their tissue-specific distribution suggests that they serve distinct biological functions in different cell types. As with CARMA1, the CARD domain of CARMA2 has been shown to specifically interact with BCL10, a protein known to function as a positive regulator of cell apoptosis and NF-kB activation. When expressed in cells, this protein activated NF-kB and induced the phosphorylation of BCL10. Alternative splicing of CARMA2 results in isoforms that possess differential effects on NF-kB activation and endoplasmic reticulum stress-induced cell death.
Synonyms:	CARMA2 Antibody, PRP, PSS1, BIMP2, CARMA2, PSORS2, Caspase recruitment domain-containing protein 14, CARD-containing MAGUK protein 2, Carma 2
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	CARD14
Reactivity:	Human
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-CARMA2 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 14 amino acid synthetic peptide near the C-terminus of human CARMA2.
Purity/Specificity:	Anti-CARMA2 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. CARMA2 antibody is human specific. At least three isoforms of CARMA2 are known to exist; this antibody will only detect isoform 1. CARMA2 antibody is predicted not to cross-react with other CARMA proteins.
Relevant Links:	• UniProtKB - Q9BXL6 • GeneID - 79092 • NCBI - NP_077015

Application Details

Tested Applications:	ELISA, IHC
Application Note:	Anti-CARMA2 Antibody has been tested for use in ELISA and Immunohistochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 113 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	User Optimized
IHC:	5 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunohistochemistry

Immunohistochemistry of CARMA2 antibody. Tissue: human placenta tissue. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: CARMA2 antibody at 5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: CARMA2 is cytoplasmic. Staining: CARMA2 is stained with Toluidine blue.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.