

Datasheet for 600-401-AK4**CIDE-A Antibody****Overview**

Description:	Anti-CIDE-A (RABBIT) Antibody - 600-401-AK4
Item No.:	600-401-AK4
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Mouse
Host Species:	Rabbit

Product Details

Background:	Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. These death signals finally cause the degradation of chromosomal DNA by activated DNase. DFF45/ICAD has been identified as inhibitor of caspase activated DNase DFF40/CAD. DFF45 related proteins CIDE-A and CIDE-B (for cell death-inducing DFF-like effector A and B) were recently identified. CIDE contains a new type of domain termed CIDE-N, which has high homology with the regulatory domains of DFF45/ICAD and DFF40/CAD. Expression of CIDE-A induces DNA fragmentation and activates apoptosis, which is inhibited by DFF45. CIDE-A is a DFF45-inhibitable effector that promotes cell death and DNA fragmentation. CIDE-A is expressed in many tissues.
Synonyms:	CIDE-A Antibody, CIDE-A, Cell death activator CIDE-A, Cell death-inducing DFFA-like effector A
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	CIDEA
Reactivity:	Mouse
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-CIDE-A antibody was prepared from whole rabbit serum produced by repeated immunizations with a 18 amino acid peptide near the C-terminus of human CIDE-A.
Purity/Specificity:	Anti-CIDE-A Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. It has no cross activity to CIDE-B.
Relevant Links:	• UniProtKB - O60543 • GeneID - 1149 • NCBI - AF041378

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-CIDE-A Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 25 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000
IF:	20 µg/mL
IHC:	2.5 µg/mL
WB:	0.5 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Wet Ice
----------------------------	---------

Storage Condition: Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.


Expiration: Expiration date is one (1) year from date of receipt.

Images




Immunofluorescence Microscopy

Immunofluorescence Microscopy of CIDE-A antibody.
Tissue: Mouse heart cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: CIDE-A antibody at 20 µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: CIDE-A is nuclear and is also localized in lipid droplets. Staining: CIDE-A as red fluorescent signal.



Immunohistochemistry

Immunohistochemistry of CIDE-A antibody. Tissue: Mouse heart tissue. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: CIDE-A antibody at 5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: CIDE-A is nuclear and is also localized in lipid droplets. Staining: CIDE-A as precipitated brown signal with hematoxylin blue nuclear counterstain.

**Western Blot**

Western Blot of CIDE-A antibody. Lane 1: Mouse heart tissue lysate. Load: 35 µg per lane. Primary antibody: CIDE-A antibody at 1:500 for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 24.7 kDa, 27 kDa for CIDE-A

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.