

**Datasheet for 600-401-AT6****DcR1 Antibody****Overview**

|                      |   |
|----------------------|---|
| <b>Description:</b>  | Anti-DcR1 (RABBIT) Antibody - 600-401-AT6 |
| <b>Item No.:</b>     | 600-401-AT6                               |
| <b>Size:</b>         | 100 µg                                    |
| <b>Applications:</b> | ELISA, IF, WB                             |
| <b>Reactivity:</b>   | Human, Mouse, Rat                         |
| <b>Host Species:</b> | Rabbit                                    |

**Product Details**

|                      |  |
|----------------------|--|
| <b>Background:</b>   | Apoptosis is induced by certain cytokines including TNF and Fas ligand in the TNF family through their death domain containing receptors. TRAIL/Apo2L is a new member of the TNF family and induces apoptosis of a variety of tumor cell lines. DR4 and DR5 are the recently identified functional receptors for TRAIL. Two decoy receptors for TRAIL have been identified and designated DcR1/TRID/TRAIL-R3/LIT and DcR2/TRAIL-R4/TRUNDD. DcR1 has extracellular TRAIL-binding domain but lacks intracellular signaling domain. It is a glycosphospholipid-anchored cell surface protein. DcR1 transcripts are expressed in many normal human tissues but not in most cancer cell lines. Overexpression of DcR1 did not induce apoptosis, but attenuated TRAIL-induced apoptosis. |
| <b>Synonyms:</b>     | DcR1 Antibody, LIT, DCR1, TRID, CD263, TRAILR3, TRAIL-R3, DCR1-TNFR, LIT, UNQ321/PRO366, Tumor necrosis factor receptor superfamily member 10C, Decoy TRAIL receptor without death domain, DcR1  |
| <b>Host Species:</b> | Rabbit   |
| <b>Clonality:</b>    | Polyclonal   |
| <b>Format:</b>       | IgG  |

**Target Details**

|                        |                    |
|------------------------|--------------------|
| <b>Gene Name:</b>      | TNFRSF10C          |
| <b>Reactivity:</b>     | Human, Mouse, Rat  |
| <b>Immunogen Type:</b> | Conjugated Peptide |

|                            |   |
|----------------------------|---|
| <b>Immunogen:</b>          | Anti-DcR1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a peptide corresponding to amino acids in the extracellular domain of human DcR1 precursor. |
| <b>Purity/Specificity:</b> | Anti-DcR1 Antibody is DEAE purified. Cross reactivity with DcR1 from other sources has not been determined.   |
| <b>Relevant Links:</b>     | <ul style="list-style-type: none"><li>• <a href="#">UniProtKB - O14798</a></li><li>• <a href="#">GeneID - 8794</a></li><li>• <a href="#">NCBI - AF012536</a></li></ul>                      |

## Application Details

|                             |  |
|-----------------------------|--|
| <b>Tested Applications:</b> | ELISA, IF, WB  |
| <b>Application Note:</b>    | Anti-DcR1 Antibody has been tested for use in ELISA, Western Blotting and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 27 kDa in Western Blots of specific cell lysates and tissues. |
| <b>Assay Dilutions:</b>     | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.  |
| <b>ELISA:</b>               | 1:10,000   |
| <b>IF:</b>                  | 10 µg/mL   |
| <b>WB:</b>                  | 1 µg/mL  |

## Formulation


|                        |   |
|------------------------|---|
| <b>Physical State:</b> | Liquid (sterile filtered)                               |
| <b>Concentration:</b>  | 1 mg/mL by UV absorbance at 280 nm                      |
| <b>Buffer:</b>         | 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2 |
| <b>Preservative:</b>   | 0.02% (w/v) Sodium Azide                                |
| <b>Stabilizer:</b>     | None  |

## Shipping & Handling

|                            |  |
|----------------------------|--|
| <b>Shipping Condition:</b> | Wet Ice  |
| <b>Storage Condition:</b>  | Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures. |


**Expiration:** Expiration date is one (1) year from date of receipt.

## Images




### Immunofluorescence Microscopy

Immunofluorescence Microscopy of DcR1 antibody. Tissue: Rat liver tissue. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: DcR1 antibody at 10  $\mu\text{g}/\text{mL}$  for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: DcR1 is localized in the cell membrane in lipid-anchors. Staining: DcR1 as red fluorescent signal with blue and green counterstains.



### Immunofluorescence Microscopy

Immunofluorescence Microscopy of DcR1 antibody. Tissue: Rat liver cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: DcR1 antibody at 20  $\mu\text{g}/\text{mL}$  for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: DcR1 is localized in the cell membrane in lipid-anchors. Staining: DcR1 as red fluorescent signal.

**Western Blot**

Western Blot of DcR1 antibody. Lane 1: HeLa cell lysate. Lane 2: Mouse liver tissue lysate. Lane 3: Rat liver tissue lysate. Load: 35 µg per lane. Primary antibody: DcR1 antibody at 1 µg/mL for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 27.4 kDa, 66 kDa for DcR1.

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.