

Datasheet for 600-401-B28S**Muscle Glycogen Synthase Phospho S641 Antibody****Overview**

Description:	Anti-Muscle Glycogen Synthase pS641 (RABBIT) Antibody - 600-401-B28S
Item No.:	600-401-B28S
Size:	25 µL
Applications:	ELISA, IHC, WB
Reactivity:	Human, Mouse
Host Species:	Rabbit

Product Details

Background:	Anti-Glycogen Synthase pS641 antibody is validated by IHC, Western Blot and ELISA. Human muscle glycogen synthase (GS) is responsible for the biosynthesis of glycogen from phosphorylated glucose units. Mammalian liver and muscle contain GS consisting of four subunits with a total molecular weight of 360,000. GS is subject to regulation through both allosteric and covalent modification and occurs in two forms: the phosphorylated inactive form, and the dephosphorylated active form. GS is inactivated by the serine/threonine kinase called glycogen synthase kinase-3 β that mainly functions to phosphorylate muscle glycogen synthase. This antibody is specific for the phosphorylated form of GS at S641. Phosphorylation of GS at S641 has been associated with Antiphospholipid Antibody Syndrome.
Synonyms:	rabbit anti-muscle glycogen synthase pS641 antibody, rabbit anti-glycogen synthase pS641 antibody, Glycogen antibody, Glycogen synthase 1 (muscle) antibody, Glycogen synthase 1 antibody, Glycogen synthase1 antibody, GYS 1 antibody, GYS-1 antibody, GYS1 antibody, Starch synthase muscle antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	GYS1
Reactivity:	Human, Mouse

PTM Specificity:	Phosphorylation
Immunogen Type:	Conjugated Peptide
Immunogen:	Human Muscle Glycogen Synthase phospho peptide corresponding to a region of the human protein conjugated to Keyhole Limpet Hemocyanin (KLH).
Purity/Specificity:	Phospho Glycogen Synthase pS641 antibody is directed against human muscle glycogen synthase. The product was affinity purified from monospecific antiserum by immunoaffinity purification. Antiserum was first purified against the phosphorylated form of the immunizing peptide. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. This phospho specific polyclonal antibody is specific for phosphorylated pS641 of human muscle glycogen synthase. Reactivity with non-phosphorylated human muscle glycogen synthase is less than 1% by ELISA. Cross reactivity with muscle glycogen synthase occurs in mouse tissue. Reactivity with muscle glycogen synthase from other sources has not been determined.
Relevant Links:	• UniProtKB - P13807 • NCBI - NP_001155059.1 • GeneID - 2997

Application Details

Tested Applications:	ELISA, IHC, WB
Application Note:	This phospho specific polyclonal antibody was tested by immunohistochemistry, immunoblotting, and ELISA. By ELISA the antibody was found to be reactive with the phosphorylated form of the immunizing peptide and minimally reactive with the non-phosphorylated form of the immunizing peptide. Immunoblotting will detect human and mouse muscle glycogen synthase. Although not tested, this antibody is likely functional in immunoprecipitation.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000 - 1:25,000
IHC:	1:1,000 - 1:5,000
WB:	1:500 - 1:3,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: None

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store Phospho GSY1 antibody at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Affinity Purified Phospho-specific pS641 antibody to human muscle Glycogen Synthase (GS). Lane 1: mouse cardiac myocyte lysate mock treated. Lane 2: mouse cardiac myocyte lysate insulin treated at 10nM for 15'. Lane 3: mouse cardiac myocyte lysate insulin treated at 100nM for 15'. Lane 4: mouse cardiac myocyte lysate insulin treated at 1nM for 15'. Lane 5: mouse cardiac myocyte lysate mock treated. Lane 6: mouse cardiac myocyte lysate CLA treated at 4nM for 45'. Lane 7: mouse cardiac myocyte lysate CLA treated at 20nM for 45'. Lane 8: mouse cardiac myocyte lysate CLA treated at 100nM for 45'. Load: 12µL. Primary Antibody: pS641 at 1:1000. Secondary Antibody: HRP conjugated Gt-a-Rabbit IgG (611-103-122) at 1:5,000 dilution preceded color development using Amersham's substrate system. Other detection methods will yield similar results.




Immunohistochemistry

Immunohistochemistry with Anti-Glycogen Synthase antibody. Tissue: Human Prostate. Fixation: formalin-fixed, paraffin-embedded tissue. Antigen retrieval: heat-induced. Primary antibody: 5 µg/ml. Staining: antibody as precipitated red signal with a hematoxylin purple nuclear counterstain.

Pathway

Diagram of glycogen synthase as a component of insulin signal transduction pathways.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.