

Datasheet for 600-401-BG5**FOXO1 Antibody****Overview**

Description:	Anti-FOXO1 (RABBIT) Antibody - 600-401-BG5
Item No.:	600-401-BG5
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	FOXO1, also known as FKHR or ForkHead in Rhabdomyosarcoma, is a 70 kDa protein which is a ubiquitously expressed member of a subfamily of the forkhead homeotic gene family of transcription factors and shuttles between the cytoplasm and nucleus. FOXO transcription factors are key players of cell fate decisions, metabolism, stress resistance, tumor suppression and are regulated by growth factors, oxidative stress or nutrient deprivation. In insulin-responsive tissues, stress or nutrient abundance triggers phosphorylation by PKB/AKT, blocking nuclear translocation and activity. Genetic mutations involving FOXO1A are a cause of alveolar rhabdomyosarcoma (RMS2). Recent studies link the anti-tumor activity of FOXO1 and the process of autophagy.
Synonyms:	FOXO1 Antibody, FKH1, FKHR, FOXO1A, Forkhead box protein O1, Forkhead box protein O1A
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	FOXO1
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-FOXO1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 15 amino acid synthetic peptide near the N-terminus of human FOXO1.

Purity/Specificity: Anti-FOXO1 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with FOXO1 from other sources has not been determined.

Relevant Links:

- [UniProtKB - Q12778](#)
- [GeneID - 2308](#)
- [NCBI - NP_002006](#)

Application Details

Tested Applications: ELISA, IF, IHC, WB

Application Note: Anti-FOXO1 Antibody has been tested for use in ELISA, Western Blotting, Immunocytochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 70 kDa in Western Blots of specific cell lysates and tissues.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:10,000

IF: 20 µg/mL

WB: 1 µg/mL

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1 mg/mL by UV absorbance at 280 nm

Buffer: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2

Preservative: 0.02% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Wet Ice

Storage Condition: Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunohistochemistry

Immunocytochemistry of FOXO1 antibody. Tissue: HeLa cells. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: FOXO1 antibody at 4 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: FOXO1 is a Nuclear and cytoplasmic. Staining: FOXO1 as precipitated blue signal with pink nuclear counterstain.



Western Blot

Western Blot of FOXO1 antibody. Lane 1: HeLa cell lysate with FOXO1 antibody at 1 µg/mL. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 70 kDa, 72 kDa for FOXO1. Other band(s): FOXO1 splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.