

Datasheet for 600-401-BJ7**GATA3 Antibody****Overview**

Description:	Anti-GATA3 (RABBIT) Antibody - 600-401-BJ7
Item No.:	600-401-BJ7
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	GATA3 is a zinc finger transcription factor that was first recognized as a possible determinant of T cell development. Later studies demonstrated that it is expressed in trophoblast giant cells and regulates trophoblast-specific genes during development. Specifically, GATA3 is selectively expressed in the trophectoderm of the peri-implantation embryo and regulates CDX2, which is required for the transcriptional repression of the POU5F1/Oct4 and NANOG genes and is thus essential for the segregation of the inner cell mass and trophectoderm at the blastocyst stage. Decreased or null-expression expression of GATA3 has also been suggested to play a major role in the development and progression of luminal breast cancer.
Synonyms:	GATA3 Antibody, HDR, HDRS, Trans-acting T-cell-specific transcription factor GATA-3, GATA-binding factor 3
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	GATA3
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-GATA3 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 17 amino acid synthetic peptide near the C-terminus of human GATA3.

Purity/Specificity: Anti-GATA3 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with GATA3 from other sources has not been determined.

Relevant Links:

- [UniProtKB - P23771](#)
- [GeneID - 2625](#)
- [NCBI - NP_001002295](#)

Application Details

Tested Applications: ELISA, IF, IHC, WB

Application Note: Anti-GATA3 Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 48 kDa in Western Blots of specific cell lysates and tissues.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:10,000 - 1:20,000

IF: 20 µg/mL

IHC: 2.5 µg/mL

WB: 1-2 µg/mL

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1 mg/mL by UV absorbance at 280 nm

Buffer: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2

Preservative: 0.02% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice


Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration:


Expiration date is one (1) year from date of receipt.

Images**Immunofluorescence Microscopy**

Immunofluorescence Microscopy of GATA3 antibody.
Tissue: Rat brain cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: GATA3 antibody at 20 µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: GATA3 is nuclear. Staining: GATA3 as red fluorescent signal.

**Western Blot**

Western Blot of GATA3 antibody. Lane 1: Human brain tissue lysate with GATA3 antibody at 1 µg/mL. Lane 2: Human brain tissue lysate with GATA3 antibody at 2 µg/mL. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 48 kDa, 40 kDa for GATA3. Other band(s): GATA3 splice variants and isoforms.

**Immunohistochemistry**

Immunohistochemistry of GATA3 antibody. Tissue: Rat brain tissue. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: GATA3 antibody at 2.5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: GATA3 is nuclear. Staining: GATA3 as precipitated brown signal with blue nuclear counterstain.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.