

Datasheet for 600-401-BL8**GLS2 Antibody****Overview**

Description:	Anti-GLS2 (RABBIT) Antibody - 600-401-BL8
Item No.:	600-401-BL8
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	Phosphate-activated glutaminase, also known as Glutaminase 2 (GLS2), was initially isolated from rat liver, although it has been shown to be expressed in other tissues. Like the functionally similar, larger kidney glutaminase, GLS2 catalyzes the hydrolysis of glutamine to stoichiometric amounts of glutamate and ammonia. Expression of GLS2 is increased by p53 under both stressed and nonstressed conditions, resulting in increased levels of glutamate and alpha-ketoglutarate, which in turn results in enhanced mitochondrial respiration and ATP generation. GLS2 also regulates antioxidant defense function in cells by increasing reduced glutathione levels and decreasing ROS-levels, suggesting that GLS2 acts as a mediator of p53's role in antioxidant defense in addition to its role in energy metabolism.
Synonyms:	GLS2 Antibody, GA, GLS, LGA, hLGA, GA, Glutaminase liver isoform, mitochondrial, L-glutaminase
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	GLS2
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-GLS2 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 18 amino acid synthetic peptide near the internal region terminus of human GLS2.
Purity/Specificity:	Anti-GLS2 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. GLS2 antibody is predicted to not cross-react with other catenin family members. Multiple isoforms of GLS2 are known to exist.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - Q9UI32• GeneID - 27165• NCBI - NP_037399

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-GLS2 Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry-P, and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 66 kDa in Western Blots of specific cell lysates and tissues. Positive control: Human pancreas tissue, mouse liver tissue, rat brain tissue.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5000 - 1:10000
IF:	10-20 µg/mL
IHC:	0.5-2 µg/mL
WB:	0.5-2 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/ml by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Wet Ice
----------------------------	---------

Storage Condition: Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

Expiration: Expiration date is one (1) year from date of receipt.

Images




Western Blot

Western Blot Validation of GLS2.

Load: 15 µg of mouse brain lysate (lane 1) or rat brain lysate (lane 2).

Primary Antibody: GLS2 at 1 µg/mL for 1 h incubation at RT in 5% NFDN/TBST.

Secondary: Goat Anti-Rabbit IgG HRP conjugate at 1:10000 dilution.



Immunofluorescence Microscopy


Immunofluorescence Validation of GLS2.

Tissue: Mouse Brain.

Fixation: 4% paraformaldehyde-fixed.

Primary Antibody: GLS2 at 20 µg/mL.

Secondary: goat anti-rabbit IgG antibody at 1:500 dilution (green) and DAPI antibody (blue).

**Immunohistochemistry**

Immunohistochemistry Validation of GLS2.


Tissue: Human Liver.

Fixation: paraffin-embedded, formaldehyde and blocked with 10% serum for 1 h at RT.

Antigen retrieval: heat mediation with a citrate buffer (pH6).

Primary Antibody: anti-GLS2 antibody at 1 μ g/ml overnight at 4°C.

Secondary: goat anti-rabbit IgG H&L (HRP) at 1:250. Counter stained with Hematoxylin.

**Immunohistochemistry**

Immunohistochemistry Validation of GLS2.


Tissue: Mouse Liver.

Fixation: paraffin-embedded, formaldehyde and blocked with 10% serum for 1 h at RT.

Antigen retrieval: heat mediation with a citrate buffer (pH6).

Primary Antibody: anti-GLS2 antibody at 1 μ g/ml overnight at 4°C.

Secondary: goat anti-rabbit IgG H&L (HRP) at 1:250. Counter stained with Hematoxylin.

**Immunohistochemistry**

Immunohistochemistry Validation of GLS2.


Tissue: Rat Brain.

Fixation: paraffin-embedded, formaldehyde and blocked with 10% serum for 1 h at RT.

Antigen retrieval: heat mediation with a citrate buffer (pH6).

Primary Antibody: anti-GLS2 antibody at 2 μ g/ml overnight at 4°C.

Secondary: goat anti-rabbit IgG H&L (HRP) at 1:250. Counter stained with Hematoxylin.




Western Blot

Western Blot Validation of GLS2.

Load: 10 μ g of human pancreas lysate.


Primary Antibody: GLS2 at 2 μ g/mL for 1 h incubation at RT in 5% NFDN/TBST.

Secondary: Goat Anti-Rabbit IgG HRP conjugate at 1:10000 dilution.




Immunofluorescence Microscopy

Immunofluorescence Microscopy of Rabbit anti-GLS2 antibody. Tissue: Human Kidney cells. Primary antibody: GLS2 antibody at 20 μ g/mL. Secondary antibody: Fluorescein rabbit secondary antibody at 1:20,000. Localization: GLS2 is located in the mitochondrion. Staining: GLS2 as red fluorescent signal.



Immunohistochemistry

Immunohistochemistry of Rabbit anti-GLS2 antibody. Tissue: human kidney tissue. Primary antibody: GLS2 antibody at 5.0 μ g/mL. Secondary antibody: Rabbit secondary antibody at 1:5,000. Localization: GLS2 is located in the mitochondrion. Staining: GLS2 as precipitated purple signal.

**Western Blot**

Western Blot of Rabbit anti-GLS2 antibody. Lane A: rat kidney tissue lysate at 0.5 µg/mL. Lane B: rat kidney tissue lysate at 1µg/mL. Primary antibody: GLS2 antibody overnight at 4°C. Secondary antibody: anti-Rabbit HRP secondary antibody. Block: 5% BLOTTO. Predicted/Observed size: 66 kDa for GLS2 antibody.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.