

Datasheet for 600-401-BW4**IL-31 Antibody****Overview**

Description:	Anti-IL-31 (RABBIT) Antibody - 600-401-BW4
Item No.:	600-401-BW4
Size:	100 µg
Applications:	ELISA, IHC, WB
Reactivity:	Mouse
Host Species:	Rabbit

Product Details

Background:	Interleukin-31 (IL-31) is a recently discovered T-cell cytokine closely related to IL-6 type cytokines and is preferentially produced by T helper type 2 cells. IL-31 activity is mediated through the ligand-induced oligomerization of a dimeric receptor complex containing IL-31 receptor A and oncostatin M receptor. In response to IL-31 binding, these proteins activate the JAK/STAT and the AKT signaling pathways. RNA levels of IL-31 receptor A and oncostatin M receptor are induced in activated monocytes but are expressed constitutively in epithelial cells. IL-31, when overexpressed in transgenic mice, results in the development of pruritis, alopecia, and skin lesions and in humans may result in atopic dermatitis, suggesting that IL-31 may represent a novel target for antipruritic drug development.
Synonyms:	IL-31 Antibody, 1700013B14Rik, Interleukin-31, IL-31
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	IL31
Reactivity:	Mouse
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-IL-31 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 16 amino acid synthetic peptide from near the N-terminus of mouse IL-31.

Purity/Specificity: Anti-IL-31 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with IL-31 from other sources has not been determined.

Relevant Links:

- [UniProtKB - Q6EAL8](#)
- [GeneID - 76399](#)
- [NCBI - AAS82846](#)

Application Details

Tested Applications: ELISA, IHC, WB

Application Note: Anti-IL-31 Antibody has been tested for use in ELISA, Western Blotting, and Immunohistochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 18 kDa in Western Blots of specific cell lysates and tissues.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:25,000 - 1:100,000

IHC: 10 µg/mL

WB: 2.5-10 µg/mL

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1.0 mg/ml by UV absorbance at 280 nm

Buffer: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2

Preservative: 0.02% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunohistochemistry

Immunohistochemistry of IL-31 antibody. Tissue: rat skeletal muscle tissue. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: IL-31 antibody at 10 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: IL-31 is secreted. Staining: IL-31 is stained with hematoxylin purple nuclear counterstain.



Western Blot

Western Blot of IL-31 antibody in RAW264.7 cell lysate. Lane A: IL-31 antibody at 2.5 µg/mL. Lane B: IL-31 antibody at 5 µg/mL. Lane C: IL-31 antibody at 10 µg/mL. Load: 35 µg per lane. Primary antibody: IL-31 antibody at designated concentrations for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 18 kDa, 31 kDa for IL-31. Other band(s): IL-31 splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.