

Datasheet for 600-401-BZ5**IRAK2 Antibody****Overview**

Description:	Anti-IRAK2 (RABBIT) Antibody - 600-401-BZ5
Item No.:	600-401-BZ5
Size:	100 µg
Applications:	ELISA, IF, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	The pro-inflammatory cytokine IL-1 induces cellular response through two subunits of its receptor, IL-1 receptor I (IL-1RI) and IL-1 receptor accessory protein (IL-1RAcP). IL-1 receptor-associated kinase (IRAK) mediates activation of NF-κB, which is a pivotal transcription factor mediating inflammatory and immune response. A novel member in the IRAK/Pelle family was recently identified and designated IRAK2. Both IRAK and IRAK2 recruit to the subunits of the IL-1R complex after IL-1 binding and lead to NF-κB activation. IRAKs also associate with Toll-like receptor (TLR) and the dominant negative mutants of IRAKs inhibit LPS-induced NF-κB activation. Members in IRAK/Pelle family play a central role in IL-1R and TLR mediated inflammatory responses to cytokine IL-1 and LPS. IRAK2 is expressed in a variety of human tissues.
Synonyms:	IRAK-2 Antibody, IRAK-2, Interleukin-1 receptor-associated kinase-like 2
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	IRAK2
Reactivity:	Human
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-IRAK2 antibody was prepared from whole rabbit serum produced by repeated immunizations with a peptide corresponding to amino acids near the C-terminus of human IRAK2.
Purity/Specificity:	Anti-IRAK-2 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Anti-IRAK2 has no cross response to IRAK.
Relevant Links:	• UniProtKB - O43187 • GeneID - 3656 • NCBI - AF026273

Application Details

Tested Applications:	ELISA, IF, WB
Application Note:	Anti-IRAK-2 Antibody has been tested for use in ELISA, Western Blotting and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 69 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000
IF:	10 µg/mL
WB:	.5 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunofluorescence Microscopy

Immunofluorescence Microscopy of IRK-2 antibody.


Tissue: HeLa cells. Fixation: 0.5% PFA. Antigen retrieval:

not required. Primary antibody: IRK-2 antibody at 10

µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT.

Localization: IRK-2 is nuclear, cytoplasmic, and is also localized in the plasma and endoplasmic membranes.

Staining: IRK-2 as red fluorescent signal.



Immunofluorescence Microscopy

Immunofluorescence Microscopy of IRK-2 antibody.


Tissue: HeLa cells. Fixation: 0.5% PFA. Antigen retrieval:

not required. Primary antibody: IRK-2 antibody at 20

µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT.

Localization: IRK-2 is nuclear, cytoplasmic, and is also localized in the plasma and endoplasmic membranes.

Staining: IRK-2 as red fluorescent signal.

**Western Blot**

Western Blot of IRAK-2 antibody. Lane 1: K562 whole cell lysate. Load: 35 µg per lane. Primary antibody: IRAK-2 antibody at 1:500 for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 69.4 kDa, 54 kDa for IRAK-2.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.