

Datasheet for 600-401-CZ6**NDST2 Antibody****Overview**

Description:	Anti-NDST2 (RABBIT) Antibody - 600-401-CZ6
Item No.:	600-401-CZ6
Size:	100 µg
Applications:	ELISA, IF, WB
Reactivity:	Human, Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	The N-deacetylase/N-sulfotransferase (heparan glucosaminyl) 2 (NDST2) protein, also known as N-heparan sulfate sulfotransferase 2, and the highly related protein NDST1 have key roles in the sulfation of heparan sulfate proteoglycans, molecules that are present of cell surfaces and the extracellular matrix (1,2). NDST1 and NDST2 are dispensable for mesodermal differentiation into osteoblasts but necessary for induction of adipocytes and neural cells (3).
Synonyms:	NDST2 Antibody, NST2, HSST2, N-heparan sulfate sulfotransferase 2, NDST-2
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	NDST2
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-NDST2 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 16 amino acid peptide near the N-terminus of human NDST2

Purity/Specificity: Anti-NDST2 antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. NDST2 antibody is human, mouse and rat reactive. At least two isoforms of NDST2 are known to exist; this antibody will only detect the larger isoform. NDST2 antibody is predicted not to cross-react with other members of the NDST protein family.

Relevant Links:

- [UniProtKB - P52849](#)
- [GeneID - 8509](#)
- [NCBI - NP_003626](#)

Application Details

Tested Applications: ELISA, IF, WB

Application Note: Anti-NDST2 Antibody has been tested for use in ELISA, Western Blotting and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 101 kDa in Western Blots of specific cell lysates and tissues.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:10,000 - 1:20,000

IF: 20 µg/mL

WB: 1-2 µg/mL

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1 mg/mL by UV absorbance at 280 nm

Buffer: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2

Preservative: 0.02% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunofluorescence Microscopy

Immunofluorescence Microscopy of Rabbit anti-NDST2 antibody. Tissue: Daudi cells. Primary antibody: NDST2 antibody at 20 $\mu\text{g}/\text{mL}$. Secondary antibody: Fluorescein rabbit secondary antibody at 1:20,000. Localization: NDST2 is located on the membrane. Staining: NDST2 as red fluorescent signal.



Western Blot

Western Blot of Rabbit anti-NDST2 antibody. Lane A: A-20 cell lysate in the absence of blocking peptide. Lane B: A-20 cell lysate in presence of blocking peptide. Primary antibody: NDST2 antibody at 1 $\mu\text{g}/\text{mL}$ overnight at 4 $^{\circ}\text{C}$. Secondary antibody: Goat anti-Rabbit HRP secondary antibody. Block: 5% BLOTTO. Predicted/Observed size: 97 kDa, 101 kDa for NDST2.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.