

Datasheet for 600-401-DA2**Neuritin Antibody****Overview**

Description:	Anti-Neuritin (RABBIT) Antibody - 600-401-DA2
Item No.:	600-401-DA2
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse
Host Species:	Rabbit

Product Details

Background:	As the nervous system of a complex organism develops, it establishes functional networks through the growth and retraction of synaptic connections from growing axons and dendrites. This synaptic remodeling involves neuro-transmitter signaling, activation of neurotrophin receptors and alterations in gene expression. One such gene whose expression is increased by neural activity is neuritin, a GPI-anchored protein that is expressed in postmitotic differentiating neurons of the developing nervous system. Its expression is also induced by the neurotrophins BDNF and NT-3. Purified recombinant neuritin promotes neurite outgrowth and arborization in primary embryonic neuronal cultures, suggesting that neuritin may play a role as a downstream effector of activity-induced neurite outgrowth. More recent experiments have shown that neuritin is required for the androgen-induced axonal elongation in motor neurons and is upregulated following spinal cord injury, suggesting that neuritin may also play a role in survival and axonal regeneration.
Synonyms:	Neuritin Antibody, NRN, dJ380B8.2, NRN, Neuritin
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	NRN1
Reactivity:	Human, Mouse

Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-Neuritin antibody was prepared from whole rabbit serum produced by repeated immunizations with a 20 amino acid synthetic peptide from near the internal region of human neuritin.
Purity/Specificity:	Anti-Neuritin Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with Neuritin from other sources has not been determined.
Relevant Links:	• UniProtKB - Q9NPD7 • GeneID - 51299 • NCBI - NP_057672

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-Neuritin Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 15 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:50,000 - 1:100,000
IF:	20 µg/mL
IHC:	2.5 µg/mL
WB:	5 and 10 µg/mL

Formulation


Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------


Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images




Immunohistochemistry

Immunohistochemistry of Neutrin antibody. Tissue: Mouse heart tissue. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: Neutrin antibody at 5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: Neutrin is nuclear and cytoplasmic. Staining: Neutrin as precipitated red signal with hematoxylin purple nuclear counterstain.



Immunofluorescence Microscopy

Immunofluorescence Microscopy of Neutrin antibody. Tissue: Mouse heart cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: Neutrin antibody at 20 µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Staining: Neutrin as red fluorescent signal.

**Western Blot**

Western Blot of Neutrin antibody. Lane A: Daudi cell lysate at 5 µg/mL. Lane B: Daudi cell lysate at 10 µg/mL. Load: 35 µg per lane. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 15.3 kDa, ~15 kDa for Neutrin. Other band(s): Neutrin splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.