

Datasheet for 600-401-DC4**NKX2-4 Antibody****Overview**

Description:	Anti-NKX2-4 (RABBIT) Antibody - 600-401-DC4
Item No.:	600-401-DC4
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse
Host Species:	Rabbit

Product Details

Background:	NKX2-4 (NK2 homeobox 4) is a member of a family of transcription factors that are involved in embryonic development and cell fate. NKX2-4 is closely related to NKX2-1 and NKX2-4 and is specifically localized to the ventral diencephalon. Recent experiments in <i>Xenopus</i> have suggested that the homeodomain protein XANF is necessary for the activation of NKX2-4, but does not directly regulate its expression.
Synonyms:	NKX2-4 Antibody, NKX2D, NKX2.4, NKX2D, Homeobox protein NK-2 homolog D
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	NKX2-4
Reactivity:	Human, Mouse
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-NKX2-4 antibody was prepared from whole rabbit serum produced by repeated immunizations with an 18 amino acid synthetic peptide near the internal region of human NKX2-4.

Purity/Specificity:	Anti-NKX2-4 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. NKX2-4 antibody is predicted to not cross-react with other NK2 homeobox family members. At least two isoforms of NKX2-4 are known to exist; this antibody will detect both isoforms.
Relevant Links:	• UniProtKB - Q9H2Z4 • GeneID - 644524 • NCBI - NP_149416

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-NKX2-4 Antibody has been tested for use in ELISA, Western Blotting, Immunocytochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 36 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000
IF:	20 µg/mL
IHC:	2.5 µg/mL
WB:	1 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunofluorescence Microscopy

Immunofluorescence of NKX2-4.

Tissue: A20 cells.

Primary Antibody: NKX2-4 antibody at 20 µg/mL.




Immunocytochemistry

Immunocytochemistry of NKX2-4.

Cell: A20 cells.

Primary Antibody: NKX2-4 antibody at 2.5 µg/mL.

**Western Blot**

Western blot analysis of NKX2-4.

Load: A20 cell lysate.

Primary Antibody: NKX2-4 antibody at 1 μ g/mL in (A) the absence and (B) the presence of blocking peptide.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.