

Datasheet for 600-401-DJ9**PARK2 Antibody****Overview**

Description:	Anti-PARK2 (RABBIT) Antibody - 600-401-DJ9
Item No.:	600-401-DJ9
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Parkinson's disease (PD) is a neurodegenerative disease whose symptoms include tremors, rigidity, bradykinesia, and postural instability (1). Mutations in the PARK2 gene are known to cause Parkinson disease and autosomal recessive juvenile Parkinson disease (2). The PARK2 protein is a component of a multiprotein E3 ubiquitin ligase complex that mediates the targeting of substrate proteins for proteasomal degradation (3). Recent studies have suggested that PARK2 expression reduces the mitochondrial accumulation of the apoptosis protein Bax under basal conditions and directly ubiquitinates Bax, thereby promoting cell survival (4).
Synonyms:	rabbit anti-Parkin pS101 Antibody, E3 ubiquitin-protein ligase parkin, Parkin, Parkin RBR E3 ubiquitin-protein ligase, Parkinson juvenile disease protein 2, Parkinson disease protein 2, PRKN, PARK2
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	PRKN
Reactivity:	Human
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-PARK2 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 17 amino acid peptide near the N-terminus of human PARK2

Purity/Specificity: Anti-PARK2 antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. PARK2 antibody is human specific.

Relevant Links:

- [UniProtKB - O60260](#)
- [GeneID - 5071](#)
- [NCBI - NP_004553](#)

Application Details

Tested Applications: ELISA, IF, IHC, WB

Application Note: Anti-PARK2 Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 52 kDa in Western Blots of specific cell lysates and tissues.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:10,000-1:20,000

IF: 20 µg/mL

IHC: 5 µg/mL

WB: 1-2 µg/mL

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2

Preservative: 0.02% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images




Immunohistochemistry

Immunohistochemistry of Rabbit anti-PARK2 antibody. Tissue: human brain. Primary antibody: PARK2 antibody at 5 $\mu\text{g}/\text{mL}$. Secondary antibody: Peroxidase rabbit secondary antibody at 1:5,000. Localization: PARK2 is cytoplasmic and nuclear. Staining: PARK2 as precipitated brown signal.



Immunofluorescence Microscopy

Immunofluorescence Microscopy of Rabbit anti-PARK2 antibody. Tissue: human brain. Primary antibody: PARK2 antibody at 20 $\mu\text{g}/\text{mL}$. Secondary antibody: Fluorescein rabbit secondary antibody at 1:20,000. Localization: PARK2 is cytoplasmic and nuclear. Staining: PARK2 as red fluorescent signal.

**Western Blot**

Western Blot of Rabbit anti-PARK2 antibody. Lane A: human cerebellum tissue lysate. Primary antibody: PARK2 antibody at 1 μ g/mL overnight at 4°C. Secondary antibody: Goat anti-Rabbit HRP secondary antibody. Block: 5% BLOTTO. Predicted/Observed size: 51 kDa, 55 kDa for PARK2.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.