

**Datasheet for 600-401-DL8****PHAP I Antibody****Overview**

<b>Description:</b>	Anti-PHAP I (RABBIT) Antibody - 600-401-DL8
<b>Item No.:</b>	600-401-DL8
<b>Size:</b>	100 µg
<b>Applications:</b>	ELISA, IF, IHC, WB
<b>Reactivity:</b>	Human, Mouse, Rat
<b>Host Species:</b>	Rabbit

**Product Details**

<b>Background:</b>	Apoptosis is related to many diseases and development. Caspase-9 plays a central role in cell death induced by a variety of apoptosis activators. Cytochrome c, after released from mitochondria, binds to Apaf-1, which forms an apoptosome that in turn binds to and activate procaspase-9. Activated caspase-9 cleaves and activates the effector caspases (caspase-3, -6 and -7), which are responsible for the proteolytic cleavage of many key proteins in apoptosis. The tumor suppressor putative HLA-DR-associated proteins (PHAPs) were recently identified as important regulators of mitochondrion apoptosis. PHAP appears to facilitate apoptosome-mediated caspase-9 activation and to stimulate the mitochondrial apoptotic pathway. PHAP was also shown to oppose both Ras- and Myc-mediated cell transformation.
<b>Synonyms:</b>	PHAP I Antibody, LANP, MAPM, PP32, HPPCn, PHAP1, PHAPI, I1PP2A, C15orf1, LANP, Acidic leucine-rich nuclear phosphoprotein 32 family member A, Acidic nuclear phosphoprotein pp32
<b>Host Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	IgG

**Target Details**

<b>Gene Name:</b>	ANP32A
<b>Reactivity:</b>	Human, Mouse, Rat
<b>Immunogen Type:</b>	Conjugated Peptide

<b>Immunogen:</b>	Anti-PHAP I antibody was raised with a synthetic peptide corresponding to amino acids at C-terminus of human PHAP I.
<b>Purity/Specificity:</b>	Anti-PHAP I Antibody is DEAE purified. This polyclonal antibody has no cross-reaction to PHAP I2a and PHAP III.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P39687</a></li><li>• <a href="#">GeneID - 8125</a></li><li>• <a href="#">NCBI - P39687</a></li></ul>

## Application Details

<b>Tested Applications:</b>	ELISA, IF, IHC, WB
<b>Application Note:</b>	Anti-PHAP I Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 29 kDa in Western Blots of specific cell lysates and tissues.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:10,000
<b>IF:</b>	20 µg/mL
<b>IHC:</b>	2 µg/mL
<b>WB:</b>	1 µg/mL

## Formulation

<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	1 mg/mL by UV absorbance at 280 nm
<b>Buffer:</b>	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.02% (w/v) Sodium Azide
<b>Stabilizer:</b>	None


## Shipping & Handling

<b>Shipping Condition:</b>	Wet Ice
----------------------------	---------

**Storage Condition:** Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.


**Expiration:** Expiration date is one (1) year from date of receipt.

## Images




### Immunohistochemistry

Immunohistochemistry of PHAP I antibody. Tissue: mouse small intestine tissue. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: PHAP I antibody at 2 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Staining: PHAP I is stained with hematoxylin purple nuclear counterstain.



### Immunofluorescence Microscopy

Immunofluorescence Microscopy of PHAP I antibody. Cell Type: mouse small intestine cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: PHAP I antibody at 20 µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: PHAP I is nuclear and cytoplasmic. Staining: PHAP I as red fluorescent signal.

**Western Blot**

Western Blot of PHAP I antibody. Lane A: human Raji cell. Lane B: mouse testis tissue lysates. Lane C: rat testis tissue lysates. Load: 35  $\mu$ g per lane. Primary antibody: PHAP I antibody at 1  $\mu$ g/ml for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 20 kDa, 32 kDa for PHAP I. Other band(s): PHAP I splice variants and isoforms.

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.