

Datasheet for 600-401-FE6**TIM-1 Antibody****Overview**

Description:	Anti-TIM-1 (RABBIT) Antibody - 600-401-FE6
Item No.:	600-401-FE6
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse
Host Species:	Rabbit

Product Details

Background:	The human form of TIM-1 was initially discovered as a membrane glycoprotein through which the hepatitis A virus can gain entry into a cell. It was also identified as kidney injury molecule 1 (Kim-1), a predicted adhesion molecule that is upregulated on the surfaces of kidney epithelia. It is also expressed on T helper 2 (Th2) cells of the immune system, and following the binding of its natural ligand TIM-4, stimulates T cell expansion and cytokine production in response to viral challenge. It has been suggested that hyperactivation of TIM-1 leads to an increased level of Th2 responsiveness and asthma susceptibility, and antibodies to TIM-1 may therefore be a novel approach to treating asthma.
Synonyms:	TIM-1 Antibody, TIM, KIM1, TIM1, HAVCR, KIM-1, TIMD1, TIMD-1, HAVCR-1, Hepatitis A virus cellular receptor 1, Kidney injury molecule 1, HAVcr-1, T-cell immunoglobulin and mucin domain-containing protein 1, T-cell immunoglobulin mucin receptor 1, T-cell membrane protein 1, CD365
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	HAVCR1
Reactivity:	Human, Mouse
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-TIM-1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 16 amino acid synthetic peptide from near the N-terminus of human TIM-1.
Purity/Specificity:	Anti-TIM-1 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with TIM-1 from other sources has not been determined.
Relevant Links:	• UniProtKB - Q96D42 • GeneID - 26762 • NCBI - NP_001166864.1

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-TIM-1 Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 39 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:20,000
IF:	20 µg/mL
IHC:	10 µg/mL
WB:	1-2 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/ml by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Wet Ice
----------------------------	---------

Storage Condition: Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunofluorescence Microscopy


Immunofluorescence of TIM-1.

Tissue: Human Testis.

Fixation: 4% paraformaldehyde-fixed.

Primary Antibody: TIM-1 at 10 µg/mL.

Secondary: goat anti-rabbit IgG secondary antibody at 1:500 dilution (red) and DAPI staining (blue).



Immunofluorescence Microscopy


Immunofluorescence of TIM-1.

Tissue: Mouse Kidney.

Fixation: 4% paraformaldehyde-fixed.

Primary Antibody: TIM-1 at 10 µg/mL.

Secondary: goat anti-rabbit IgG secondary antibody at 1:500 dilution (red) and DAPI staining (blue).




Western Blot

Western Blot of TIM-1.

Load: 15 µg of human cell lysates per lane.

Primary Antibody: TIM-1 at 8 µg/mL for overnight incubation at 4° C in 5% NFDm/TBST.

Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.




Western Blot

Western Blot of TIM-1.

Load: 15 µg of lysates per lane.

Primary Antibody: Top: TIM-1 (p/n 600-401-FE6) at 8 µg/mL, Middle: TIM1 at 1 µg/mL for overnight incubation at 4° C; Bottom: beta-actin at 1 µg/mL and GAPDH at 0.02 µg/mL for 1h incubation at RT, in 5% NFDm/TBST.

Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.




Western Blot

Western Blot of TIM-1.

Load: 15 µg of cell lysates per lane. Lane 1: HeLa cells were transfected with control siRNAs, Lane 2: TIM-1 siRNA Knockdown in HeLa cells.

Primary Antibody: TIM-1 at 8 µg/mL for 1 h incubation at RT in 5% NFDm/TBST.

Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.




Western Blot

Western Blot of TIM-1.

Load: 15 μ g of mouse tissue lysates.

Primary Antibody: TIM-1 at 2 μ g/mL for 1h incubation at RT in 5% NFDM/TBST.

Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.




Immunohistochemistry

Immunohistochemistry of TIM-1 antibody. Tissue: human uterus tissue. Fixation: formalin fixed paraffin embedded.

Antigen retrieval: not required. Primary antibody: TIM-1 antibody at 10 μ g/mL for 1 h at RT. Secondary antibody:

Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: TIM-1 is nuclear and chromosomal.

Staining: TIM-1 is stained with toluidine blue.




Immunofluorescence Microscopy

Immunofluorescence Microscopy of TIM-1 antibody. Cell

Type: human uterus cells. Fixation: 0.5% PFA. Antigen

retrieval: not required. Primary antibody: TIM-1 antibody at 20 μ g/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT.

Localization: TIM-1 is nuclear and chromosomal. Staining: TIM-1 as red fluorescent signal.



Western Blot

Western Blot of TIM-1 antibody in human uterus tissue lysate. Lane A: TIM-1 antibody at 1 μ g/mL. Lane B: TIM-1 antibody at 2 μ g/mL. Load: 35 μ g per lane. Primary antibody: TIM-1 antibody at designated concentrations for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 75 kDa, 48 kDa for TIM-1. Other band(s): TIM-1 splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.