

**Datasheet for 600-401-FG6****TLR9 Antibody****Overview**

<b>Description:</b>	Anti-TLR9 (RABBIT) Antibody - 600-401-FG6
<b>Item No.:</b>	600-401-FG6
<b>Size:</b>	100 µg
<b>Applications:</b>	ELISA, IF, IHC, WB
<b>Reactivity:</b>	Human, Mouse
<b>Host Species:</b>	Rabbit

**Product Details**

<b>Background:</b>	Toll-like receptors (TLRs) are evolutionarily conserved pattern-recognition molecules resembling the toll proteins that mediate antimicrobial responses in <i>Drosophila</i> . These proteins recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. TLR9 forms a subfamily along with TLR7 and TLR8 that recognize viral RNA and CpG DNA sequences and are localized in intracellular acidic compartments such as the phagolysosome. Unlike other TLRs which act through adaptor molecules such as TOLLIP, TIRAP, TRIF, and MyD88 to activate various kinases and transcription factors to respond to potential infection, TLR9 is strictly dependent on MyD88.
<b>Synonyms:</b>	TLR9 Antibody, CD289, Toll-like receptor 9
<b>Host Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	IgG

**Target Details**

<b>Gene Name:</b>	TLR9
<b>Reactivity:</b>	Human, Mouse
<b>Immunogen Type:</b>	Conjugated Peptide
<b>Immunogen:</b>	Anti-TLR9 antibody was prepared from whole rabbit serum produced by repeated immunizations with a peptide corresponding to 16 amino acids near the C-terminus of human TLR9.

**Purity/Specificity:** Anti-TLR9 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with TLR9 from other sources has not been determined.

**Relevant Links:**

- [UniProtKB - Q9NR96](#)
- [GeneID - 54106](#)
- [NCBI - NP\\_059138.1](#)

## Application Details

**Tested Applications:** ELISA, IF, IHC, WB

**Application Note:** Anti-TLR9 Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 116 kDa in Western Blots of specific cell lysates and tissues.

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

**ELISA:** 0.5-2 µg/mL

**IF:** 10 µg/mL

**IHC:** 2 µg/mL

**WB:** 0.5-2 µg/mL

## Formulation

**Physical State:** Liquid (sterile filtered)

**Concentration:** 1 mg/mL by UV absorbance at 280 nm

**Buffer:** 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2

**Preservative:** 0.02% (w/v) Sodium Azide

**Stabilizer:** None


## Shipping & Handling

**Shipping Condition:** Dry Ice

**Storage Condition:** Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


**Expiration:** Expiration date is one (1) year from date of receipt.

## Images




### Immunohistochemistry

Immunohistochemistry of TLR9 antibody. Cell Type: Mouse Spleen cells. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: TLR9 antibody at 2  $\mu\text{g}/\text{mL}$  for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: TLR9 is located in the cell membrane. Staining: TLR9 is stained with toluidine blue.



### Immunofluorescence Microscopy

Immunofluorescence Microscopy of TLR9 antibody. Cell Type: Mouse Spleen cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: TLR9 antibody at 10  $\mu\text{g}/\text{mL}$  for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: TLR9 is located in the cell membrane. Staining: TLR9 as red fluorescent signal.

**Western Blot**

Western Blot of TLR9 antibody in mouse spleen cell lysate. Lane A: TLR9 antibody at 0.5  $\mu\text{g}/\text{mL}$ . Lane B: TLR9 antibody at 1  $\mu\text{g}/\text{mL}$ . Lane C: TLR9 antibody at 2  $\mu\text{g}/\text{mL}$ . Load: 35  $\mu\text{g}$  per lane. Primary antibody: TLR9 antibody at designated concentrations for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 96 kDa, 94 kDa for TLR9. Other band(s): TLR9 splice variants and isoforms.

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.