

## Datasheet for 600-401-GH6

**STAT3 (Internal) Antibody****Overview**

<b>Description:</b>	Anti-STAT3 (RABBIT) Antibody - 600-401-GH6
<b>Item No.:</b>	600-401-GH6
<b>Size:</b>	100 µg
<b>Applications:</b>	Dot Blot
<b>Reactivity:</b>	Human
<b>Host Species:</b>	Rabbit

**Product Details**

**Background:** Signal transducer and activator of transcription 3 (Stat3) belongs to a family of cytoplasmic transcription factors that can be activated by phosphorylation by its cell surface receptor. Stat3 plays a key role in many cellular processes such as cell growth and apoptosis. It also mediates cellular responses to interleukins, KITLG/SCF, EGF, IFN-alpha and other growth factors and may mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Stat3 forms a homodimer or a heterodimer with a related family member (e.g. STAT1). Activation occurs through phosphorylation of tyrosine 705 and serine 727. Phosphorylation of Stat3 at Tyr705 induces Stat3 dimerization and nuclear translocation. Serine phosphorylation is important for stable DNA-binding of Stat3 homodimers and maximal transcriptional activity. Stat3 can have a dual role in cancer, it has been found that Stat3 protein can promote oncogenesis and have a tumor suppressor role depending upon the mutational background of the tumor.

**Synonyms:** rabbit anti-STAT3 antibody, STAT-3, STAT 3, Signal transducer and activator of transcription 3, Acute-phase response factor, APRF

**Host Species:** Rabbit

**Clonality:** Polyclonal

**Format:** IgG

**Target Details**

**Gene Name:** STAT3

**Reactivity:** Human

<b>Immunogen Type:</b>	Conjugated Peptide
<b>Immunogen:</b>	STAT3 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with synthetic monomethylated or asymmetric dimethylated peptides surrounding Arginine 31 of human STAT.
<b>Purity/Specificity:</b>	Anti-STAT3 was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with all STAT3 modifications. Cross-reactivity is seen against STAT1 and STAT4. BLAST analysis was used to suggest cross-reactivity with Human, mouse, and rat. Cross-reactivity with STAT3 from other sources has not been determined.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li><a href="#">UniProtKB - P40763</a></li></ul>

## Application Details

<b>Tested Applications:</b>	Dot Blot
<b>Application Note:</b>	Anti-STAT3 antibody has been tested by Dot Blot and is useful for Western Blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~88 kDa corresponding to STAT3 protein by Western Blotting in the appropriate cell lysate or extract.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:5000
<b>WB:</b>	1:500
<b>Other:</b>	Dot Blot: 1:1000

## Formulation

<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	1.0 mg/ml by UV absorbance at 280 nm
<b>Buffer:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.05% (w/v) Sodium Azide
<b>Stabilizer:</b>	30% Glycerol


## Shipping & Handling

<b>Shipping Condition:</b>	Dry Ice
----------------------------	---------

**Storage Condition:** Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

## Images



### Dot Blot

Dot blot of Rabbit Anti-STAT3 antibody. Antigen: non-modified, monomethylated and asymmetric dimethylated forms of the immunizing peptide from the internal portion of the STAT3 protein. Load: 100, 10, or 1 picomolar as indicated. Primary antibody: STAT3 (Internal) antibody at 1:1000 for 90 min at RT. (Date: 10/22/14; Exposure time: 3.6 seconds).

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.