

Datasheet for 600-401-GN3**Glut2 Antibody****Overview**

Description:	Anti-Glut2 (RABBIT) Antibody - 600-401-GN3
Item No.:	600-401-GN3
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse
Host Species:	Rabbit

Product Details

Background: The Anti-Glut2 antibody was designed, produced, and validated as part of the Joy Cappel Young Investigator Award (JCYIA). The glucose transporter GLUT2 is a transmembrane carrier protein that allows protein facilitated glucose movement across cell membranes. GLUT2 is expressed in the plasma membranes of the liver, intestine, renal tubular cells, pancreatic islet beta cells, as well as in the portal and hypothalamic areas. Due to its low affinity and high capacity, GLUT2 transports dietary sugars, glucose, galactose and fructose in high concentrations, displaying large bidirectional fluxes in and out of cells. In pancreatic beta cells, GLUT2 is essential for glucose-stimulated insulin secretion. GLUT2 expression is necessary for the physiological control of glucose-sensitive genes, and its inactivation in the liver leads to impaired glucose-stimulated insulin secretion. In the nervous system, GLUT2-dependent glucose sensing regulates feeding, thermoregulation and pancreatic islet cell mass and function, as well as sympathetic and parasympathetic activities. In humans, inactivating mutations in GLUT2 cause Fanconi–Bickel syndrome, which is characterized by hepatomegaly and kidney disease. Anti-Glut2 is ideal for researchers interested in studying glucose transport mediated by Glut2 protein in the fields of diabetes, obesity, metabolism, and neuroscience research.

Synonyms:	rabbit anti-Glut2 antibody, Solute carrier family 2 facilitated glucose transporter member 2, Glucose transporter type 2 liver, GLUT-2 antibody, Slc2a2, Glut 2 Antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	Slc2a2
Reactivity:	Human, Mouse
Immunogen Type:	Conjugated Peptide
Immunogen:	This affinity-purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to the C-terminal domain of mouse Glut2 protein.
Purity/Specificity:	This affinity-purified Glut-2 antibody is directed against Glut2 protein. The product was affinity purified from monospecific antiserum by immunoaffinity purification. A BLAST analysis was used to suggest cross-reactivity with Glut2 from mouse and rat based on a 100% homology with the immunizing sequence. Reactivity against homologues from other sources is not known.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P14246• NCBI - NP_112474.2• GeneID - 20526

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-Glut-2 antibody has been tested for use in ELISA, immunohistochemistry, immunofluorescence, and by western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 57.1 kDa in size corresponding to Glut2 protein by western blotting in the appropriate stimulated tissue or cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:30,000-1:90:000
FLISA:	1
IHC:	4µg/mL
WB:	1:500-1:1000


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None



Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images





A: Negative Control
B: Glut2 staining in human kidney (20x)
C: Glut2 staining in human kidney (40x)




Immunohistochemistry

Immunohistochemistry with anti-Glut2 antibody showing Glut2 staining in nucleus and cytoplasm of ductal epithelium and of renal glomeruli in human kidney at 20x and 40x (B & C). Formalin fixed/paraffin embedded sections were subjected to heat induced epitope retrieval (HIER) at pH 6.2 and then incubated with rabbit anti-mouse Glut2 antibody at 4.0 µg/ml for 60 minutes. The reaction was developed using MACH 1 universal HRP polymer detection system and visualized with 3'3-diamino-benzidine substrate (DAB).




Immunohistochemistry

Immunohistochemistry of Anti-Glut2 Antibody. Tissue: mouse kidney. Antigen retrieval: Heat Induced, slides incubated in sodium citrate buffer for 1hr at 90°C. Primary: Rabbit Anti-Glut2 Antibody at 5µg/mL overnight. Blocking: 2% goat serum in TBST. Secondary: Alexa 488 at 1µg/mL for 2hrs at room temperature.

**Western Blot**

Western Blot of Rabbit anti-Glut2 antibody. Lane 1: HEK293 lysate (p/n W09-000-365). Load: 10 μ g per lane. Primary antibody: Glut2 antibody at 1:1000 for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody (p/n 611-103-122) at 1:40,000 for 30 min at RT. Block: Blocking Buffer (p/n MB-070) overnight at 4°C. Predicted/Observed size: ~60 kDa.

**Western Blot**

Western Blot of Rabbit Anti-Glut2 Antibody. Lane 1: Molecular Weight. Lane 2: Mouse Kidney WCL (p/n W10-000-T014). Lane 3: MEF WCL (p/n W10-001-371). Lane 4: 3T3 WCL (p/n W10-000-358). Lane 5: HeLa WCL (p/n W09-000-364). Lane 6: HEK293 WCL (p/n W09-000-365). Load: 20 μ g per pane. Primary antibody: Anti-Glut2 at 1 μ g/mL overnight at 2-8°C. Secondary Antibody: Goat anti-Rabbit HRP (p/n 611-103-122) 1:40,000 diluted with MB-073 for 30 minutes at RT. Blocked with (p/n MB-073) Buffer. Expect: ~57.1 kDa. Exposure: 30 sec.

References

- Bathina S et al. Normal β -Cell Glut2 Expression Is not Required for Regulating Glucose-Stimulated Insulin Secretion and Systemic Glucose Homeostasis in Mice. *Biomolecules*. (2023)
- Fujino M et al. Transcription factor c-Maf deletion improves streptozotocin-induced diabetic nephropathy by directly regulating Sglt2 and Glut2. *JCI Insight*. (2023)
- Laskowska AK et al. Fruits of Hippophaë rhamnoides in human leukocytes and Caco-2 cell monolayer models-A question about their preventive role in lipopolysaccharide leakage and cytokine secretion in endotoxemia. *Front Pharmacol*. (2022)
- de Souza Cordeiro LM et al. Loss of function of renal Glut2 reverses hyperglycaemia and normalises body weight in mouse models of diabetes and obesity. *Diabetologia*. (2022)
- Liang et al. Organic cation transporter 1 (OCT1) modulates multiple cardiometabolic traits through effects on hepatic thiamine content. *PLOS Biology* (2018)
- Chhabra et al. Reduced renal sympathetic nerve activity contributes to elevated glycosuria and improved glucose tolerance in hypothalamus-specific Pomc knockout mice. *Molecular Metabolism* (2017)
- Chhabra et al. Hypothalamic POMC Deficiency Improves Glucose Tolerance Despite Insulin Resistance by Increasing Glycosuria. *Diabetes* (2016)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.