

Datasheet for 600-401-I52**Histone H3 R2me1 Antibody****Overview**

Description:	Anti-Histone H3 [Monomethyl Arg2] (RABBIT) Antibody - 600-401-I52
Item No.:	600-401-I52
Size:	50 µg
Applications:	Dot Blot, WB
Reactivity:	Human, C. elegans
Host Species:	Rabbit

Product Details

Background:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the telomeric copy. Anti-Histone H3 are ideal for researchers interested in Chromatin Modifiers, Chromatin Research, Histones and Modified Histones, and Epigenetics research.
Synonyms:	rabbit anti-Histone H3 monomethyl Arg2 antibody, H3.3B, H3 histone, family 3A, H3.3AH3F3H3F3B, histone H3.3, MGC87783, MGC87782, H3R2me1
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	HIST2H3C
Reactivity:	Human, C. elegans

PTM Specificity:	Methylation
Immunogen Type:	Conjugated Peptide
Immunogen:	Histone H3 [Monomethyl Arg2] affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic monomethylated peptide surrounding Arginine 2 of human Histone H3.2.
Purity/Specificity:	Anti-Histone H3 [Monomethyl Arg2] was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with human Histone H3.2. A BLAST analysis was used to suggest cross-reactivity with Human, mouse, and C. elegans. Predicted to react with many species including rat, chicken, Xenopus, Drosophila, and plant based on 100% sequence homology. Cross-reactivity with Histone H3 from other sources has not been determined.
Relevant Links:	• UniProtKB - Q71DI3 • NCBI - NP_001005464 • GeneID - 126961

Application Details

Tested Applications:	Dot Blot, WB
Application Note:	Anti-Histone H3 [Monomethyl Arg2] antibody is tested for Western Blot and Dot Blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~15.4 kDa corresponding to Histone H3 protein by Western Blotting in HeLa histone prep lysate or the appropriate cell lysate or extract. Epi-Plus™ antibody production in collaboration with Novus Biologicals.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
WB:	1:500


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.71 mg/ml by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling


Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Dot Blot

Dot Blot of Rabbit Anti-Histone H3 R3 Me1 Antibody. Lane 1: R2 NM. Lane 2: R2 Me1. Lane 3: R2 Me2s. Lane 4: R2 Me2a. Load: 100, 10, 1 picomoles of peptide. Primary antibody: Histone H3 R3 Me1 antibody at 1:1,000 for 45 min at 4°C. Secondary Antibody: Dylight™488 rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C.



Western Blot

Western Blot of Rabbit Anti-Histone H3 [Monomethyl Arg2] Antibody. Lane 1: HeLa histone prep lysate. Load: 30 µg per lane. Primary antibody: Histone H3 [Monomethyl Arg2] at 1:500 for overnight at 4°C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: ~15 kDa. Other band(s): None.

**Western Blot**

Western Blot of Rabbit Anti-Histone H3 [Monomethyl Arg2] Antibody. Lane 1: *C. elegans* embryo lysate. Load: 30 µg per lane. Primary antibody: Histone H3 [Monomethyl Arg2] at 1:500 for overnight at 4°C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: ~15 kDa. Other band(s): None.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.