

Datasheet for 600-401-J69**HDAC5 (internal) Antibody****Overview**

Description:	Anti-HDAC5 (RABBIT) Antibody - 600-401-J69
Item No.:	600-401-J69
Size:	100 µg
Applications:	ELISA, IHC, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	HDAC5 is a ubiquitous protein responsible for the deacetylation of lysine residues on the N-terminal part of the core histones. A tag for epigenetic repression, Anti-HDAC5 is an enzyme involved in transcriptional regulation, cell cycle progression and developmental events. HDAC5 is a class II histone deacetylase as defined by its similarities to other members of this class. HDAC5 is a major player in muscle differentiation, acting by repressing transcription of myocyte enhancer MEF2C. During differentiation it moves from the nucleus into the cytoplasm, allowing for myocyte enhancer factor expression. Anti-HDAC5 antibody is ideal for researchers interested in Stem Cell , Cardiovascular , or Epigenetics and Nuclear Signaling research.
Synonyms:	rabbit anti-HDAC-5 antibody, HDAC5, HDAC 5, HD 5 antibody, histone deacetylase 5 antibody, Histone deacetylase-5, HD5 antibody, NY-CO-9
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	HDAC5
Reactivity:	Human
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-HDAC5 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to the internal region surrounding 325-375aa of human HDAC5 protein.
Purity/Specificity:	Anti-HDAC5 was affinity purified from monospecific antiserum by immunoaffinity chromatography. A BLAST analysis was used to suggest cross-reactivity with human and hamster based on 100% sequence homology. Cross-reactivity with HDAC5 from other sources has not been determined.
Relevant Links:	• UniProtKB - Q9UQL6

Application Details

Tested Applications:	ELISA, IHC, WB
Application Note:	Anti-HDAC5 antibody tested in ELISA, Immunohistochemistry, and Western Blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~122kDa corresponding to the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1: 10,000
IHC:	5 µg/mL
WB:	1ug/ml


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.1 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer:	30% Glycerol

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Immunohistochemistry

Immunohistochemistry of Rabbit anti-HDAC5 antibody.
Tissue: Colon. Fixation: formalin fixed paraffin embedded.
Antigen retrieval: not required. Primary antibody: HDAC5 antibody at 5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Staining: HDAC5 as precipitated red signal with hematoxylin purple nuclear counterstain.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.