Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 600-401-MV9

Glucagon Antibody

Overview

Description:	Anti-Glucagon (RABBIT) Antibody - 600-401-MV9
Item No.:	600-401-MV9
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Glucagon is a member of a multigene family that includes secretin. Glucagon is a 29-amino acid pancreatic hormone that counteracts the glucose-lowering action of insulin by stimulating glycogenolysis and gluconeogenesis. The human glucagon gene is approximately 9.4 kb long, contains 6 exons and 5 introns, and assigned to 2q36-2q37. This antibody is suitable for researchers interested in metabolic diseases, like diabetes, cell proliferation, differentiation, apoptosis, GCPR signaling, and calcium signaling research.
Synonyms:	Rabbit Anti-Glucagon, Pro-glucagon, Glicentin, Oxyntomodulin, OXM, OXY, Glucagon
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	GCG
Reactivity:	Human
Immunogen Type:	Conjugated Peptide
Immunogen:	Glucagon antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal portion of human Glucagon.

Purity/Specificity:	This affinity purified antibody is directed against human Glucagon. This product was affinity purified from monospecific antiserum by immunoaffinity purification. A BLAST analysis was used to suggest cross-reactivity with the antigen based on 100% homology with the immunizing sequence to human, chimpanzee, and bonobo.
Relevant Links:	• GeneID - 2641 • UniProtKB - P01275 • NCBI - NP_002045.1

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-Glucagon is tested in ELISA, IF, IHC-P, and Western Blot. Expect a band approximately ~20.9 kDa corresponding to the appropriate cell lysate or extract. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:24,800-1:44,800
IF:	5-15µg/mL
IHC:	User Optimized
WB:	User Optimized

Formulation


Physical State:	Liquid (sterile filtered)
Concentration:	1.18 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling


Shipping Condition:	Dry Ice

Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.


Images



Western Blot
Western Blot of Rabbit Anti-Glucagon Antibody.
Lane 1: Opal Prestained Molecular Weight (p/n MB-210-0500).
Lane 2: COS-7 Lysate - reduced (20µg).
Lane 3: BSA Conjugated Glucagon peptide - reduced (0.02µg).
Lane 4: Insulin - reduced (0.05µg).
Primary Antibody: Anti-Glucagon [Rabbit] Antibody at 1.0µg/mL overnight at 2-8°C.
Secondary Antibody: Goat Anti-Rabbit IgG (MX10) Peroxidase conjugated at 1:70,000 for 30mins at RT.
Block: Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) for 1hr at RT.
Expected MW: ~21kDa. Observed MW: endogenous detection in COS-7 Lysate at ~35kDa. Glucagon peptide is detected at the MW of BSA. No cross-reactivity with insulin is observed.
Exposure: 25 sec.



ELISA
ELISA Results of Rabbit Anti-Glucagon Antibody.
Each well was coated with 1µg of conjugate. The starting concentration of antibody in the dilution series was 5 µg/ml. The titer is 1:34,800 Glucagon - Free peptide [Green Line], 1:47200 Glucagon Standard - BSA conjugated [Blue Line], and 1:20,500 Glucagon - BSA conjugated [Purple Line]. Each point on the Y-axis represents a 3-fold dilution. 3% Fish Gel (p/n MB-066-0100), HRP conjugated Goat anti-Rabbit IgG (H&L) (p/n 611-1302), and TMB substrate (p/n TMB-1000) were used for detection.



Immunohistochemistry
Immunohistochemistry results using Rabbit Anti-Glucagon Antibody.
Tissue: alpha cells in CD1 mouse pancreatic islets.
Fixation: 4% paraformaldehyde.
Antigen Retrieval: 10mM Sodium Citrate buffer for 10 mins at 95-100°C.
Blocking: PBS, 1% ovalbumin, 0.3% Triton X-100.
Primary Antibody: Anti-Glucagon at 1:100 overnight at RT.
Secondary Antibody: Anti-Rabbit Alexa Fluor 488 at 1:500 for 1hr at RT.
Original magnification 20x.
Independently Validated by antibodies-online GmbH (ABIN7448121). Courtesy of Prof. Merighi, University of Turin.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.