

Datasheet for 600-401-R05**DDT Antibody****Overview**

Description:	Anti-DDT (RABBIT) Antibody - 600-401-R05
Item No.:	600-401-R05
Size:	100 µg
Applications:	IHC, WB, IP
Reactivity:	Human, Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	DDT, D-dopachrome tautomerization, converts D-dopachrome into 5,6-dihydroxyindole. Northern blot analysis revealed that DDT was expressed as a 0.6-kb mRNA in all tissues tested, with the strongest expression in liver. The DDT gene in human and mouse is identical in exon structure to the MIF gene. Both genes have 2 introns that are located at equivalent positions, relative to a 2-fold repeat in protein structure. The genes for DDT and MIF are closely linked on human chromosome 22 and mouse chromosome 10. This antibody is suitable for researchers interested in stem cell research.
Synonyms:	D-dopachrome tautomerase, Phenylpyruvate tautomerase II
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	DDT
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide
Immunogen:	DDT affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a sequence at the C-terminal of human DDT.

Purity/Specificity: Anti-DDT antibody is directed against human DDT protein. The product was affinity purified from monospecific antiserum by immunoaffinity purification. A BLAST analysis was used to suggest reactivity with this protein from human, mouse, and rat based on homology for the immunogen sequence. Cross reactivity with DDT from other sources has not been determined.

Relevant Links:

- [UniProtKB - P30046](#)
- [GeneID - 1652](#)
- [NCBI - NP_001077861.1](#)

Application Details

Tested Applications:	IHC, WB
Suggested Applications:	IP (Based on references)
Application Note:	Anti-DDT is tested for Immunohistochemistry-P and Western Blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~12.7 kDa corresponding to the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
IHC:	1:100-1:500
WB:	0.5 µg/mL

Formulation


Physical State:	Lyophilized
Concentration:	0.5 mg/mL by UV absorbance at 280 nm
Buffer:	0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃
Preservative:	0.05mg Thimerosal
Stabilizer:	5mg BSA
Reconstitution Volume:	200µL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition: Ambient

Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.


Images



Immunohistochemistry

Immunohistochemistry analysis of DDT using anti-DDT antibody.

DDT was detected in paraffin-embedded section of human lung cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1µg/ml rabbit anti-DDT Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



Western Blot

Western blot analysis of DDT using anti-DDT antibody (from recombinant protein, Left) and anti-DDT antibody (600-401-R05, from peptide, Right).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30µg of sample under reducing conditions.

Lane 1: human hepatocellular carcinoma tumor tissue (HCCT) lysates, Lane 2: rat liver tissue lysates, Lane 3: mouse liver tissue lysates.

After Electrophoresis, proteins were transferred to a nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with either affinity purified rabbit anti-DDT rec antigen polyclonal antibody or affinity purified rabbit anti-DDT antigen polyclonal antibody (p/n 600-401-R05) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for DDT at approximately 13-15 kDa. The expected band size for DDT is at 13 kDa.

References

- Song S et al. D-dopachrome tautomerase contributes to lung epithelial repair via atypical chemokine receptor 3-dependent Akt signaling. *EBioMedicine*. (2021)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.