

Datasheet for 600-401-Y49**APP Antibody****Overview**

Description:	Anti-APP (RABBIT) Antibody - 600-401-Y49
Item No.:	600-401-Y49
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human, Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	Accumulation of the amyloid-beta peptide (Abeta) in the cerebral cortex is a critical event in the pathogenesis of Alzheimer's disease. The beta-amyloid protein precursor (APP) is cleaved by beta-secretase, producing a soluble derivative of the protein and a membrane anchored 99-amino acid carboxy-terminal fragment (C99). The C99 fragment serves as substrate for gamma-secretase to generate the 4 kDa amyloid-beta peptide (Abeta), which is deposited in the brains of all suffers of Alzheimer's disease.
Synonyms:	APP Antibody, AAA, AD1, PN2, ABPP, APPI, CVAP, ABETA, PN-II, CTFgamma, A4, Amyloid beta A4 protein, APP
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	APP
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-APP antibody was prepared from whole rabbit serum produced by repeated immunizations with a 15 amino acid peptide near the C-terminus of human APP.

Purity/Specificity: Anti-APP Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with APP from other sources has not been determined.

Relevant Links:

- [UniProtKB - P05067](#)
- [GeneID - 351](#)
- [NCBI - CAA30050](#)

Application Details

Tested Applications: ELISA, IF, IHC, WB

Application Note: Anti-APP Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Expect a band at approximately 87 kDa in Western Blots using positive control mouse and rat brain tissue, which can also be used in IHC and IF, or specific cell lysates and tissues. Specific conditions for reactivity should be optimized by the end user.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:10,000

IF: 20 µg/mL

IHC: 2 µg/mL

WB: 0.5-1 µg/mL

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2

Preservative: 0.02% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Wet Ice


Storage Condition: Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

Expiration:


Expiration date is one (1) year from date of receipt.

Images**Immunofluorescence Microscopy**

Immunofluorescence Microscopy of APP antibody. Tissue: Rat brain cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: APP antibody at 20 µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: APP is localized in the cell membrane. Staining: APP as red fluorescent signal.

**Immunohistochemistry**

Immunohistochemistry of APP antibody. Tissue: Rat brain. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: APP antibody at 2 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: APP is localized in the cell membrane. Staining: APP as precipitated orange signal with hematoxylin purple nuclear counterstain.

**Western Blot**

Western Blot of APP antibody. Lane 1: Mouse brain tissue lysate. Lane 2: Rat brain tissue lysate. Load: 35 µg per lane. Primary antibody: APP antibody at 2 µg/mL for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 86.9 kDa, 105 kDa for APP. Other band(s): APP splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.