

Datasheet for 600-401-Z26**BANF1 Antibody****Overview**

Description:	Anti-BANF1 (RABBIT) Antibody - 600-401-Z26
Item No.:	600-401-Z26
Size:	100 µg
Applications:	ELISA, IF, WB
Reactivity:	Human, Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	Barrier-to-autointegration factor 1 (BANF1) is a conserved chromatin protein that non-specifically binds double-stranded DNA. BAF also interacts with a family of nuclear proteins that include LAP2, emerin, and MAN1. It is also a host cell component of retroviral pre-integration complexes (PICs), including that of HIV. BAF will bind to p55 Gag (the structural precursor of HIV-1 virions) as well as its cleaved product matrix. In addition to being a host cell component of the PIC, it is thought that BAF is also present at low levels in incoming virions, and thus might contribute to the assembly or activity of HIV-1 PICs through direct binding to matrix as well as DNA.
Synonyms:	BANF1 Antibody, BAF, NGPS, BCRP1, D14S1460, BAF, BCRG1, Barrier-to-autointegration factor, Breakpoint cluster region protein 1
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	BANF1
Reactivity:	Human, Mouse, Rat
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-BANF1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a 19 amino acid synthetic peptide from near the N-terminus of human BANF1.
Purity/Specificity:	Anti-BANF1 Antibody was affinity purified from monospecific antiserum by immunoaffinity chromatography. Cross reactivity with BANF1 from other sources has not been determined.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - O75531• GeneID - 8815• NCBI - AAH05942

Application Details

Tested Applications:	ELISA, IF, WB
Application Note:	Anti-BANF1 Antibody has been tested for use in ELISA, Western Blotting and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 20 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5000 - 1:20,000
IF:	20 µg/mL
WB:	0.5 - 2 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images




Western Blot

Western Blot using BANF. Loading: 15 µg of lysates per lane. Antibodies: BANF1 p/n 600-401-Z25 (2 µg/mL), BANF1 p/n 600-401-Z26 (2 µg/mL) and beta-actin (1.5 µg/mL), 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.



Western Blot


Western Blot Validation. Loading: 15 µg of human cell line lysates per lane. Antibodies: BANF1, (2 µg/mL), 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.



Western Blot


Western Blot Validation in Mouse and Rat Cell Lines.

Loading: 15 µg of lysates per lane. Antibodies: BANF1 (2 µg/mL), 1h incubation at RT in 5% NFDN/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.



Immunofluorescence Microscopy

Immunofluorescence Microscopy of BANF1 antibody in human kidney tissue. Tissue: human kidney tissue. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: BANF1 antibody at 20 µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: BANF1 is nuclear and cytoplasmic. Staining: BANF1 as red fluorescent signal with bis-benzimide (blue) nuclear counterstain.



Western Blot

Western Blot of BANF1 antibody. Lane A: Mouse kidney tissue lysate at 0.5 µg/mL. Lane B: Mouse kidney tissue lysate at 1 µg/mL. Lane C: Mouse kidney tissue lysate at 2 µg/mL. Load: 35 µg per lane. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 10 kDa, ~24 kDa for BANF1.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.