

## Datasheet for 600-402-098

## Biotin Antibody Fluorescein Conjugated

### Overview

|                      |  |
|----------------------|--|
| <b>Description:</b>  | Anti-Biotin (RABBIT) Antibody Fluorescein Conjugated - 600-402-098 |
| <b>Item No.:</b>     | 600-402-098  |
| <b>Size:</b>         | 100 µL   |
| <b>Applications:</b> | WB   |
| <b>Reactivity:</b>   | Biotin   |
| <b>Host Species:</b> | Rabbit   |

### Product Details

|                      |  |
|----------------------|--|
| <b>Background:</b>   | Biotin Antibody detects Biotin. Biotin is a water-soluble B-complex vitamin (vitamin B7). It is composed of a ureido (tetrahydroimidizalone) ring fused with a tetrahydrothiophene ring. A valeric acid substituent is attached to one of the carbon atoms of the tetrahydrothiophene ring. Biotin is a coenzyme for carboxylase enzymes, involved in the synthesis of fatty acids, isoleucine, and valine, and in gluconeogenesis. Biotin is necessary for cell growth, the production of fatty acids, and the metabolism of fats and amino acids. Anti-Biotin Fluorescein Conjugated Antibody is ideal for investigators involved in Cell Signaling and Cell Biology research. |
| <b>Synonyms:</b>     | rabbit anti-Biotin Antibody fluorescein Conjugation, FITC conjugated rabbit anti-Biotin Antibody   |
| <b>Host Species:</b> | Rabbit   |
| <b>Conjugate:</b>    | Fluorescein (FITC)   |
| <b>Clonality:</b>    | Polyclonal   |
| <b>Format:</b>       | IgG  |
| <b>F/P Ratio:</b>    | 2.8  |

### Target Details

|                        |  |
|------------------------|--|
| <b>Reactivity:</b>     | Biotin   |
| <b>Immunogen Type:</b> | Other  |
| <b>Immunogen:</b>      | Biotin conjugated to Keyhole Limpet Hemocyanin (KLH) |

**Purity/Specificity:** Biotin Antibody Fluorescein Conjugated was prepared from monospecific antiserum by immunoaffinity chromatography using Biotin coupled to sepharose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Biotinylated IgG and Biotinylated Bovine Serum Albumin.

---

## Application Details

**Tested Applications:** WB

---

**Application Note:** Biotin Antibody Fluorescein Conjugated has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

---

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

---

**IF:** 1:500-1:2,500

---

## Formulation

**Physical State:** Lyophilized

---

**Concentration:** 1.0 mg/mL by UV absorbance at 280 nm

---

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

---

**Preservative:** 0.01% (w/v) Sodium Azide

---

**Stabilizer:** 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

---

**Reconstitution Volume:** 100  $\mu$ L

---

**Reconstitution Buffer:** Restore with deionized water (or equivalent)

---

## Shipping & Handling

**Shipping Condition:** Ambient

---


**Storage Condition:** Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

---

**Expiration:** Expiration date is one (1) year from date of receipt.

---

## Images



### Western Blot

Western Blot of Rabbit anti-Biotin Fluorescein Conjugated Antibody. Lane 1: Biotin Conjugated Guinea Pig IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Fluorescein rabbit secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 28 & 55 kDa, 28 & 55 kDa for Guinea Pig IgG. Other band(s): None.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.