

Datasheet for 600-403-215

GFP Antibody Peroxidase Conjugated**Overview**

Description:	Anti-GFP (RABBIT) Antibody Peroxidase Conjugated - 600-403-215
Item No.:	600-403-215
Size:	100 µg
Applications:	ELISA, WB
Reactivity:	GFP, eGFP, rGFP
Host Species:	Rabbit

Product Details

Background:	Green fluorescent protein is a 27 kDa protein produced from the jellyfish <i>Aequorea victoria</i> , which emits green light (emission peak at a wavelength of 509nm) when excited by blue light. GFP is an important tool in cell biology research. GFP is widely used enabling researchers to visualize and localize GFP-tagged proteins within living cells without the need for chemical staining.
Synonyms:	rabbit anti-GFP antibody peroxidase conjugation, HRP conjugated rabbit anti-GFP antibody, Green Fluorescent Protein, GFP antibody, Green Fluorescent Protein antibody, EGFP, enhanced Green Fluorescent Protein, <i>Aequorea victoria</i> , Jellyfish
Host Species:	Rabbit
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	GFP, eGFP, rGFP
Immunogen Type:	Recombinant Protein
Immunogen:	Anti-Green Fluorescent Protein (GFP) is produced by immunizing GFP containing fusion protein that corresponds to the full length amino acid sequence (246aa) derived from the jellyfish <i>Aequorea victoria</i> .

Purity/Specificity: GFP Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Green Fluorescent Protein (*Aequorea victoria*) coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, anti-Peroxidase and purified and partially purified Green Fluorescent Protein (*Aequorea victoria*) Serum. No reaction was observed against Human, Mouse and Rat Serum Proteins.

Relevant Links:

- [UniProtKB - P42212](#)

Application Details

Tested Applications: ELISA, WB

Application Note: Anti-GFP Antibody is designed to detect GFP and its variants. This GFP Peroxidase conjugated antibody has been tested by ELISA and Western blot. This product can be used to detect GFP by ELISA (sandwich or capture) for the direct binding of antigen and recognizes wild type, recombinant and enhanced forms of GFP. Biotin conjugated polyclonal anti-GFP used in a sandwich ELISA is well suited to titrate GFP in solution using this antibody in combination with Rockland's monoclonal anti-GFP (600-301-215) using either form of the antibody as the capture or detection antibody. However, use the monoclonal form only for the detection of wild type or recombinant GFP as this form does not sufficiently detect 'enhanced' GFP. The detection antibody is typically conjugated to biotin and subsequently reacted with streptavidin conjugated HRP (code # S000-03). Fluorochrome conjugated polyclonal anti-GFP can be used to detect GFP by immunofluorescence microscopy in prokaryotic (*E.coli*) and eukaryotic (CHO cells) expression systems and can detect GFP containing inserts. Significant amplification of signal is achieved using fluorochrome conjugated polyclonal anti-GFP relative to the fluorescence of GFP alone. For immunoblotting use either alkaline phosphatase or peroxidase conjugated polyclonal anti-GFP to detect GFP or GFP containing proteins on western blots. Optimal titers for applications should be determined by the researcher.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:10,000 - 1:50,000

IHC: 1:500 - 1:2,000

WB: 1:2,000 - 1:5,000

Formulation

Physical State: Lyophilized

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2


Preservative: 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	100 μ L
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling


Shipping Condition:	Ambient
Storage Condition:	Store GFP Antibody at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Co-IP was performed using stage-11 embryos injected with GFP-Rab11fp5 RNA (1 ng) and ephrinB1 deletion mutants (300 pg each) as indicated. Fig. 2. PMID: 33462110




Western Blot

(A) RNAs were injected into two D1 blastomeres at the eight-cell stage as indicated. Embryos were harvested at 5, 10 and 15 h after injection and western blot analysis was performed. Fig 5. PMID: 33462110

Western Blot

Western Blot of Anti-GFP Antibody Peroxidase Conjugate. Lane 1: 50ng of GFP. Lane 2: none. Primary antibody: none. Secondary antibody: Anti-GFP Antibody Peroxidase Conjugate secondary antibody was used at 1:1000 for 45 min at RT. Block: 1% BSA-TTBS (p/n MB-013, diluted to 1X) 30 min at 20°C. Predicted/Observed size: 28 kDa for GFP. Other band(s): none.



Western Blot

Western Blot of Rabbit anti-GFP Peroxidase Conjugated Antibody. Lane 1: GFP. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Peroxidase rabbit secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 28 kDa, 28 kDa for GFP. Other band(s): None.

References

- Lines M et al. The Dishevelled C-terminus interacts with the centrosomal protein Kizuna to regulate microtubule organization during ciliogenesis. *bioRxiv [Preprint]*. (2025)
- Yoon J et al. Rab11fip5 regulates telencephalon development via ephrinB1 recycling. *Development*. (2021)
- Kakraba S et al. A Novel Microtubule-Binding Drug Attenuates and Reverses Protein Aggregation in Animal Models of Alzheimer's Disease. *Frontiers in Molecular Neuroscience* (2019)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.