

**Datasheet for 601-1204****Bovine IgG F(ab')<sub>2</sub> Antibody Fluorescein Conjugated****Overview**

|                      |                                                                                     |
|----------------------|-------------------------------------------------------------------------------------|
| <b>Description:</b>  | Goat Anti-Bovine IgG F(ab') <sub>2</sub> Antibody Fluorescein Conjugated - 601-1204 |
| <b>Item No.:</b>     | 601-1204                                                                            |
| <b>Size:</b>         | 2 mg                                                                                |
| <b>Applications:</b> | Dot Blot                                                                            |
| <b>Reactivity:</b>   | Bovine                                                                              |
| <b>Host Species:</b> | Goat                                                                                |

**Product Details**

|                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|----------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Background:</b>   | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. This product possesses the F(ab') <sub>2</sub> fragment, recognized by the two F(ab) fragments yielded from the digestion of the antibody below the disulfide bond hinge region. |
| <b>Synonyms:</b>     | goat anti-Bovine IgG F(ab') <sub>2</sub> fragment fluorescein Conjugated Antibody, goat anti-Bovine IgG Fab2 fragment FITC Conjugated Antibody                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Host Species:</b> | Goat                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Specificity:</b>  | IgG F(ab') <sub>2</sub>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Conjugate:</b>    | Fluorescein (FITC)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Clonality:</b>    | Polyclonal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Format:</b>       | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>F/P Ratio:</b>    | 3.26                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |

**Target Details**

|                    |                                         |
|--------------------|-----------------------------------------|
| <b>Reactivity:</b> | Bovine                                  |
| <b>Immunogen:</b>  | Bovine IgG F(ab') <sub>2</sub> fragment |

|                            |                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Purity/Specificity:</b> | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Bovine IgG, Bovine IgG F(ab') <sub>2</sub> and Bovine Serum. No reaction was observed against Bovine IgG F(c). |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

## Application Details

|                             |                                                                                                                                                                                                                                                                                                                                                 |
|-----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Tested Applications:</b> | Dot Blot                                                                                                                                                                                                                                                                                                                                        |
| <b>Application Note:</b>    | Anti-Bovine IgG F(ab') <sub>2</sub> Fluorescein Conjugated has been tested by dot blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. |
| <b>Assay Dilutions:</b>     | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.                                                                                                                                                                                                                                                 |
| <b>FC:</b>                  | 1:500 - 1:2,500                                                                                                                                                                                                                                                                                                                                 |
| <b>FLISA:</b>               | 1:10,000 - 1:50,000                                                                                                                                                                                                                                                                                                                             |
| <b>IF:</b>                  | 1:1,000 - 1:5,000                                                                                                                                                                                                                                                                                                                               |

## Formulation

|                               |                                                         |
|-------------------------------|---------------------------------------------------------|
| <b>Physical State:</b>        | Lyophilized                                             |
| <b>Concentration:</b>         | 2.0 mg/mL by UV absorbance at 280 nm                    |
| <b>Buffer:</b>                | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| <b>Preservative:</b>          | 0.01% (w/v) Sodium Azide                                |
| <b>Stabilizer:</b>            | 10 mg/ml Polyethylene Glycol (PEG-8000)                 |
| <b>Reconstitution Volume:</b> | 1.0 mL                                                  |
| <b>Reconstitution Buffer:</b> | Restore with deionized water (or equivalent)            |

## Shipping & Handling

|                            |                                                                                                                                                                                                                                                                                                                                                 |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Shipping Condition:</b> | Ambient                                                                                                                                                                                                                                                                                                                                         |
| <b>Storage Condition:</b>  | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |

**Expiration:** Expiration date is one (1) year from date of receipt.

## Images



### Dot Blot

Dot Blot of Goat anti-Bovine IgG Fab2 Antibody Fluorescein Conjugated. Antigen: Bovine IgG. Load: Lane 1 - 100 ng Lane 2 - 33.3 ng Lane 3 - 11.1 ng Lane 4 - 3.70 ng Lane 5 - 1.23 ng. Primary antibody: n/a. Secondary antibody: Goat anti-Bovine IgG Fab2 Antibody Fluorescein Conjugated at 1:1,000 for 1 HR at RT. Block: MB-070 for 1 HR at RT.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.