

Datasheet for 607-4602**Golden Syrian Hamster IgG (H&L) Antibody Biotin Conjugated****Overview**

Description:	Rabbit Anti-Golden Syrian Hamster IgG (H&L) Antibody Biotin Conjugated - 607-4602
Item No.:	607-4602
Size:	1.5 mg
Applications:	ELISA
Reactivity:	Golden Syrian Hamster
Host Species:	Rabbit

Product Details

Background:	Anti-Golden Syrian Hamster IgG Biotin Antibody generated in rabbit detects Golden Syrian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. Anti-Golden Syrian Hamster IgG Antibody is ideal for investigators in Cancer, Immunology, and Microbiology research.
Synonyms:	Golden Syrian Hamster IgG (H&L) Antibody, Rabbit-a-Golden Syrian Hamster IgG, Golden Syrian Hamster IgG in Rabbit, Rabbit IgG Secondary Antibody, rabbit anti-hamster IgG antibody
Host Species:	Rabbit
Specificity:	IgG (H&L)
Conjugate:	Biotin
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Golden Syrian Hamster
Immunogen:	Hamster IgG whole molecule
Purity/Specificity:	Anti-Golden Syrian Hamster IgG Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Hamster IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Hamster IgG and Hamster Serum.

Application Details

Tested Applications:	ELISA
Application Note:	Anti-Golden Syrian Hamster IgG Biotin Conjugation has been tested by ELISA and is assayed against 1.0 ug of Hamster IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:15,000 to 1:60,000 of the reconstitution concentration is suggested for Anti-Golden Syrian Hamster IgG.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000 - 1:100,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:10,000


Formulation

Physical State:	Lyophilized
Concentration:	1.5 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



ELISA

ELISA Results of Purified Rabbit Anti-Golden Syrian Hamster IgG Antibody Biotin Conjugated tested against purified Golden Syrian Hamster IgG Biotin. Each well was coated in duplicate with 1.0 µg of Golden Syrian Hamster IgG (p/n 007-0102). The working dilution is 1:57,000. The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using HRP Conjugation Stabilizer (p/n MB-076), Streptavidin-HRP conjugated (p/n S000-03), and TMB substrate (p/n TMBE-1000).

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.