Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 609-4303

Human IgG Fc Antibody Peroxidase Conjugated

Overview

Description:	Rabbit Anti-Human IgG Fc Antibody Peroxidase Conjugated - 609-4303
Item No.:	609-4303
Size:	2 mg
Applications:	ELISA
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Anti-Human IgG F(c) Peroxidase Conjugated generated in rabbit detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of human IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(c) Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.
Synonyms:	rabbit Anti-Human IgG F(c) Antibody peroxidase Conjugated, rabbit Anti-Human IgG Fc fragment Antibody peroxidase Conjugated, rabbit Anti-Human IgG Fc Antibody HRP Conjugated
Host Species:	Rabbit
Specificity:	IgG Fc
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Human

Immunogen:	Human IgG F(c) fragment
Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab).

Application Details

Tested Applications:	ELISA
Application Note:	Anti-Human IgG F(c) Peroxidase Conjugated Antibody has been tested by ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:50,000
IHC:	1:500 - 1:2,500
WB:	1:1,000 - 1:5,000

Formulation

Physical State:	Lyophilized
Concentration:	2.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient

Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



ELISA
ELISA results of purified Rabbit anti-Human IgG F(c) Antibody Peroxidase Conjugated tested against purified Human IgG F (c). Each well was coated in duplicate with 1.0 µg of Human IgG F(c) (p/n 009-0103). The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% fish gelatin as blocking buffer and TMB substrate p/n TMBE-1000.

References

- Ching KH et al. Generation of human sequence antibodies in OmniChicken against the highly conserved mammalian target BDNF using three distinct discovery workflows. *Mabs.* (2026)
- Tanno C et al. Artificial transcription factor-driven bidirectional expression for sustained high-level monoclonal antibody production in CHO cells. *Science Direct.* (2026)
- Kaneko Y et al. Evaluation of cellular characteristics and genome editing responses in chicken primordial germ cell lines. *J Biosci Bioeng.* (2026)
- Vuong CN et al. Heavy chain-only antibodies with a stabilized human VH in transgenic chickens for therapeutic antibody discovery. *MAbs.* (2024)
- Wulandari DA et al. Design and validation of functionalized redox-responsive hydrogel beads for high-throughput screening of antibody-secreting mammalian cells. *J Biosci Bioeng.* (2024)
- Ying B et al. High-Level Production of scFv-Fc Antibody Using an Artificial Promoter System with Transcriptional Positive Feedback Loop of Transactivator in CHO Cells. *Cells.* (2023)
- Zheng F et al. LINE-1 vectors mediate recombinant antibody gene transfer by retrotransposition in Chinese hamster ovary cells. *Biotechnol J.* (2021)

www.rockland.com
tech@rockland.com
+1 484.791.3823

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.