

Datasheet for 609-4318**Human IgG F(ab')₂ Antibody Peroxidase Conjugated Pre-Adsorbed****Overview**

Description:	Rabbit Anti-Human IgG F(ab') ₂ Antibody Peroxidase Conjugated (Min X MOUSE Serum Proteins) - 609-4318
Item No.:	609-4318
Size:	1 mg
Applications:	ELISA
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Anti-Human IgG F(ab') ₂ Peroxidase Antibody generated in rabbit recognizes the dimeric Fab portion of the human IgG molecule. Human IgG F(ab') ₂ is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab') ₂ molecules lack the Fc portion of IgG and therefore receptors that bind human IgG F(c) will not bind human IgG F(ab') ₂ molecules. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
Synonyms:	rabbit Anti-Human IgG Fab ₂ Antibody peroxidase Conjugated, Rabbit Anti-Human IgG F(ab') ₂ Antibody peroxidase Conjugated, Rabbit Anti-Human IgG Fab ₂ Fragment Antibody HRP Conjugated
Host Species:	Rabbit
Specificity:	IgG F(ab') ₂
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Human
--------------------	-------

Immunogen:	Human IgG F(ab') ₂ fragment
Purity/Specificity:	Anti-Human IgG F(ab') ₂ antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption (s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Human IgG, Human IgG F(ab') ₂ and Human Serum. No reaction was observed against Human IgG F(c) or Mouse Serum Proteins.

Application Details

Tested Applications:	ELISA
Application Note:	Anti-Human IgG F(ab') ₂ Peroxidase Antibody has been tested by ELISA and is assayed against 1.0 ug of Human IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:8,500 to 1:72,000 of the reconstitution concentration is suggested for this product. This product is also suitable for Western Blot and Immunohistochemistry. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:50,000
IHC:	1:500 - 1:2,500
WB:	1:1,000 - 1:10,000

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition:	Ambient
----------------------------	---------

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



ELISA

ELISA results of purified Rabbit Anti-Human IgG F(ab')₂ Peroxidase Conjugated Antibody tested against purified Human IgG F(ab')₂. Each well was coated in duplicate with 1.0 µg of Human IgG F(ab')₂ (blue line). The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC₅₀ is defined as the titer of the antibody. Assay performed using TMB-1000 substrate.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.