

Datasheet for 610-103-121

Mouse IgG (H&L) Antibody Peroxidase Conjugated Pre-adsorbed

Overview

Description:	Goat Anti-Mouse IgG (H&L) Antibody Peroxidase Conjugated (Min X Bv Ch Gt GP Ham Hs Hu Rb Rt & Sh Serum Proteins) - 610-103-121
Item No.:	610-103-121
Size:	1 mg
Applications:	ELISA, WB, IHC, Other
Reactivity:	Mouse
Host Species:	Goat

Product Details

Background:	Anti-Mouse IgG Peroxidase Antibody generated in goat detects reactivity to Mouse IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
Synonyms:	Goat Anti-Mouse IgG Secondary Antibody Peroxidase Conjugated, Goat Anti-Mouse IgG Secondary Antibody HRP Conjugated, GAM-HRP, Anti-mouse secondary antibody, anti-mouse HRP antibody, horseradish peroxidase conjugated secondary antibody, anti-mouse horseradish peroxidase conjugated secondary antibody
Host Species:	Goat
Specificity:	IgG (H&L)
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

F/P Ratio: 0.387

Target Details

Reactivity:	Mouse
Immunogen:	Mouse IgG whole molecule
Purity/Specificity:	HRP secondary antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption (s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.

Application Details

Tested Applications:	ELISA, WB
Suggested Applications:	IHC, Other (Based on references)
Application Note:	Anti-Mouse IgG Peroxidase Antibody has been tested by ELISA and western blot and is ideal for western blotting, Immunohistochemistry and ELISA as well as other antibody detection methods.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:180,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:20,000


Formulation

Physical State:	Lyophilized
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store secondary antibody at 4° C prior to restoration. For extended storage aliquot antibody and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Anti-Mouse IgG HRP secondary antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Peroxidase conjugated Goat anti-Mouse IgG antibody. Lane 1: Mouse IgG. Lane 2: none. Load: 50 ng per lane. Primary antibody: none. Secondary antibody: Peroxidase mouse secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 55 kDa, 28 kDa for Mouse IgG. Other band(s): none.

References

- Li M et al. Escin Rescues Blood-brain Barrier and Inhibits NLRP3 Inflammasome-mediated Pyroptosis in Rats with Superior Sagittal Sinus Thrombosis. *Int J Med Sci.* (2025)
- Chen F et al. TGF- β 1-induced apoptosis in retinal endothelial cells is implicated in retinal vein occlusion. *Exp Eye Res.* (2025)
- Zhong H et al. The zinc finger motif in the mitochondrial large ribosomal subunit protein bL36m is essential for optimal yeast mitoribosome assembly and function. *Biochim Biophys Acta Mol Cell Res.* (2024)
- Fritz K et al. Hypercapnia Causes Injury of the Cerebral Cortex and Cognitive Deficits in Newborn Piglets. *eNeuro.* (2024)
- Akiyama H et al. Novel IgE crosslinking-induced luciferase expression method using human-rat chimeric IgE receptor-carrying mast cells. *J Immunol Methods.* (2024)
- Sang Y et al. PI4KIII β -Mediated Phosphoinositides Metabolism Regulates Function of the VTA Dopaminergic Neurons and Depression-Like Behavior. *J Neurosci.* (2024)

- Koyuncu D et al. B cells in perivascular and peribronchiolar granuloma-associated lymphoid tissue and B-cell signatures identify asymptomatic Mycobacterium tuberculosis lung infection in Diversity Outbred mice. *Infect Immun.* (2024)
- Barrera C et al. Effects on Fetal Metabolic Programming and Endocannabinoid System of a Normocaloric Diet during Pregnancy and Lactation of Female Mice with Pregestational Obesity. *Nutrients.* (2023)
- Wang W et al. Matrix stiffness regulates tumor cell intravasation through expression and ESRP1-mediated alternative splicing of MENA. *Cell Rep.* (2023)
- Schenke M et al. Human-Relevant Sensitivity of iPSC-Derived Human Motor Neurons to BoNT/A1 and B1. *Toxins (Basel).* (2021)
- Yingling CV et al. FHOD formin and SRF promote post-embryonic striated muscle growth through separate pathways in *C. elegans*. *Exp Cell Res.* (2021)
- Yang X, He Z, Zhang G, et al. Evaluation of Reactivity of Monoclonal Antibodies Against Omp25 of Brucella spp. *Front Cell Infect Microbiol.* (2020)
- Amanat F, Stadlbauer D, Strohmeier S, et al. A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat. Med.* (2020)
- SchötteIndreier D et al. Low Density Lipoprotein Receptor-Related Protein-1 (LRP1) Is Involved in the Uptake of Clostridioides difficile Toxin A and Serves as an Internalizing Receptor. *Front Cell Infect Microbiol.* (2020)
- Nishida T et al. Suppression of adipocyte differentiation by low-intensity pulsed ultrasound via inhibition of insulin signaling and promotion of CCN family protein 2. *J Cell Biochem.* (2020)
- Timon-Gomez A et al. Distinct roles of mitochondrial HIGD1A and HIGD2A in respiratory complex and supercomplex biogenesis. *Cell Rep.* (2020)
- Kim MR et al. TET2 directs mammary luminal cell differentiation and endocrine response. *Nat Commun.* (2020)
- Timon-Gomez A et al. Protocol for the Analysis of Yeast and Human Mitochondrial Respiratory Chain Complexes and Supercomplexes by Blue Native Electrophoresis. *STAR Protoc.* (2020)
- Genth et al. Difference in Mono-O-Glucosylation of Ras Subtype GTPases Between Toxin A and Toxin B From Clostridioides difficile Strain 10463 and Lethal Toxin From Clostridium sordellii Strain 6018. *Frontiers in Microbiology* (2019)
- Fan et al. Upregulation of miR-185 promotes apoptosis of the human gastric cancer cell line MGC803. *Molecular Medicine Reports* (2018)
- SchötteIndreier et al. Expression and (Lacking) Internalization of the Cell Surface Receptors of Clostridioides difficile Toxin B. *Frontiers in Microbiology* (2018)
- Brown et al. FOXP3 and miR-155 cooperate to control the invasive potential of human breast cancer cells by down regulating ZEB2 independently of ZEB1. *Oncotarget* (2018)
- Lobo-Jarne T, Nývltová E, Pérez-Pérez R, et al. Human COX7A2L Regulates Complex III Biogenesis and Promotes Supercomplex Organization Remodeling without Affecting Mitochondrial Bioenergetics. *Cell Rep.* (2018)
- Ciesielska A et al. Bis (monoacylglycero) phosphate inhibits TLR4-dependent RANTES production in macrophages. *Int J Biochem Cell Biol.* (2017)

- Shen HY et al. Antitumor activity of fucoidan in anaplastic thyroid cancer via apoptosis and anti-angiogenesis. *Mol Med Rep.* (2017)
- Langer N et al. Determination of cross-reactivity of poly-and monoclonal antibodies for synthetic cannabinoids by direct SPR and ELISA. *Forensic Sci Int.* (2017)
- Ercan et al. Neuronal CTGF/CCN2 negatively regulates myelination in a mouse model of tuberous sclerosis complex. *Journal of Experimental Medicine* (2017)
- Lampi et al. Simvastatin Ameliorates Matrix Stiffness-Mediated Endothelial Monolayer Disruption. *PLOS One* (2016)
- Schelle et al. Role of p38alpha/beta MAP Kinase in Cell Susceptibility to Clostridium sordellii Lethal Toxin and Clostridium difficile Toxin B. *Toxins* (2016)
- Dickerman et al. The protein activator of protein kinase R, PACT/RAX, negatively regulates protein kinase R during mouse anterior pituitary development. *The FEBS Journal* (2015)
- Horn P et al. Circulating microparticles carry a functional endothelial nitric oxide synthase that is decreased in patients with endothelial dysfunction. *J Am Heart Assoc.* (2012)
- LeVine H et al. Dihydroxybenzoic acid isomers differentially dissociate soluble biotinyl-A β (1-42) oligomers. *Biochemistry* (2012)
- Zhang X et al. Vaccination with different M2e epitope densities confers partial protection against H5N1 influenza A virus challenge in chickens. *Intervirology.* (2011)
- McCall LI et al. Localization and induction of the A2 virulence factor in Leishmania: evidence that A2 is a stress response protein. *Mol Microbiol.* (2010)
- Reeves TM et al. Proteolysis of submembrane cytoskeletal proteins ankyrin-G and α II-spectrin following diffuse brain injury: a role in white matter vulnerability at Nodes of Ranvier. *Brain Pathology (Zurich, Switzerland)* (2010)
- Van Komen JS et al. The polybasic juxtamembrane region of Sso1p is required for SNARE function in vivo. *Eukaryot Cell.* (2005)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.