

Datasheet for 610-1602

Mouse IgG (H&L) Antibody Biotin Conjugated**Overview**

Description:	Goat Anti-Mouse IgG (H&L) Antibody Biotin Conjugated - 610-1602
Item No.:	610-1602
Size:	2 mg
Applications:	ELISA, IHC, WB
Reactivity:	Mouse
Host Species:	Goat

Product Details

Background:	Anti-Mouse IgG (H&L) biotin conjugated antibody generated in goat detects specifically mouse IgG (H&L). This secondary conjugated antibody anti-mouse is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.
Synonyms:	Goat Anti-Mouse IgG Secondary Antibody biotin Conjugation, Goat Anti-Mouse IgG Antibody biotin Conjugated, Anti-mouse IgG secondary antibody, anti-mouse IgG biotin conjugated secondary antibody
Host Species:	Goat
Specificity:	IgG (H&L)
Conjugate:	Biotin
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Mouse
Immunogen Type:	Native Protein
Immunogen:	Anti-Mouse IgG was produced by repeated immunization with Mouse IgG whole molecule in goat.

Purity/Specificity: This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Mouse IgG and Mouse Serum.

Application Details

Tested Applications:	ELISA
Suggested Applications:	IHC, WB (Based on references)
Application Note:	Anti-Mouse IgG (H&L) biotin conjugated antibody has been tested by ELISA and is suitable for highly specific immunological methods requiring extremely low background levels, lot-to-lot consistency, high titer and specificity.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:350,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:20,000


Formulation

Physical State:	Lyophilized
Concentration:	2.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



ELISA

ELISA results of purified Goat anti-Mouse IgG Antibody Biotin Conjugated tested against purified Mouse IgG. Each well was coated in duplicate with 1.0 μg of Mouse IgG (p/n 010-0102). The starting dilution of antibody was 5 $\mu\text{g}/\text{ml}$ and the X-axis represents the Log₁₀ of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC₅₀ is defined as the titer of the antibody. Assay performed using 3% fish gelatin as blocking buffer, Streptavidin Peroxidase Conjugated p/n S000-03) and TMB substrate p/n TMBE-1000.

References

- Rahman MA et al. Loss of HIV candidate vaccine efficacy in male macaques by mucosal nanoparticle immunization rescued by V2-specific response. *Nat Commun.* (2024)
- Mueller A et al. In vitro assembly of Tobacco mosaic virus coat protein variants derived from fission yeast expression clones or plants. *J Virol Methods.* (2010)
- Rowlands JC et al. Altered mammary gland differentiation and progesterone receptor expression in rats fed soy and whey proteins. *Toxicol Sci.* (2002)
- Alam R et al. Increased MCP-1, RANTES, and MIP-1 α in bronchoalveolar lavage fluid of allergic asthmatic patients. *Am J Respir Crit Care Med.* (1996)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.